

AN ROINN OIDEACHAIS  
LEAVING CERTIFICATE EXAMINATION, 1990

1697

APPLIED MATHEMATICS – HIGHER LEVEL

FRIDAY, 22 JUNE – MORNING, 9.30 – 12.00

Six questions to be answered. All questions carry equal marks.

Mathematics Tables may be obtained from the Superintendent.

Take the value of  $g$  to be  $9.8 \text{ m/s}^2$ .

Marks may be lost if all your work is not shown or you do not indicate where a calculator has been used.

1. (a) A particle is projected vertically upwards with velocity  $u$  m/s and is at a height  $h$  after  $t_1$  and  $t_2$  seconds respectively. Prove that

$$t_1 \cdot t_2 = \frac{2h}{g}$$


- (b) A car accelerates uniformly from rest to a speed  $v$  m/s. It continues at this constant speed for  $t$  seconds and then decelerates uniformly to rest.

The average speed for the journey is  $\frac{3v}{4}$ .

- (i) Draw a speed-time graph and hence, or otherwise, prove that the time for the journey is  $2t$  seconds.
- (ii) If the car-driver had observed the speed limit of  $\frac{1}{2}v$ , find the least time the journey would have taken, assuming the same acceleration and deceleration as in (i).

2. Two straight roads intersect at an angle of  $60^\circ$ . Car  $A$  moves towards the junction with uniform speed 16 m/s, while car  $B$  moves away from the junction with uniform speed 20 m/s.


- (a) Calculate the velocity of  $A$  relative to  $B$ .
- (b) If  $A$  is 450 m and  $B$  is 200 m from the intersection at a given moment, calculate the time interval in seconds until the cars
- are nearest to each other
  - are equidistant from the intersection.



3. A particle is projected from a point  $p$ , up a plane inclined at an angle  $\tan^{-1} \frac{1}{6}$  to the horizontal. The direction of projection makes an angle  $\alpha$  with the inclined plane. (The plane of projection is vertical and contains the line of greatest slope.)
- If the particle were to strike the inclined plane horizontally at a point  $q$ , show that  $\tan \alpha = \frac{3}{19}$ .
  - If the particle were to be projected from  $p$  with the same speed but at an angle  $\tan^{-1} 3$  to the inclined plane, show that it would strike the plane at right angles at  $q$ .

OVER →

4. Two blocks  $A$  and  $B$  each of mass  $m$  kg, lie at rest on horizontal rough tables. The coefficient of friction between  $A$  and the table is  $\mu$ , and between  $B$  and its table is  $\frac{1}{4}$ . The blocks are connected by a light inextensible string which passes under a smooth movable pulley of mass  $2m$  kg.




$F = ma$   
 $mg = \mu a$

- (i) Show in a diagram the forces acting on each mass when the system is released from rest.
- (ii) If  $\mu < \frac{3}{4}$ , prove that the tension in the string is  $\frac{mg(9 + 4\mu)}{16}$
- (iii) Prove that  $A$  will not move if  $\mu > \frac{3}{4}$ .

5. State the laws governing the oblique collision of two smooth elastic spheres.

A smooth elastic sphere of mass  $6$  kg rests on a smooth horizontal table. A second smooth elastic sphere of mass  $4$  kg falls vertically on it. At the moment of impact the line of centres makes an angle of  $45^\circ$  with the vertical, and the velocity of the falling sphere is  $u$ . The  $6$  kg sphere moves horizontally after the collision.



- (i) Explain why the principle of conservation of momentum may be applied horizontally.
- (ii) Hence, or otherwise, prove that the speed of the  $6$  kg mass after impact is  $\frac{u(1 + e)}{4}$  where  $e$  is the coefficient of restitution between the two spheres.
- (iii) If  $e = \frac{1}{3}$ , prove that the loss of kinetic energy due to the impact is

$$\frac{2u^2}{3}$$


6. (a) A particle starts from rest, and moves with simple harmonic motion of period  $6n$  seconds. Show that the particle moves from the position of maximum velocity to the position in which the velocity is half the maximum in  $n$  seconds.
- (b) The depth of water in a harbour is assumed to rise and fall with time in simple harmonic motion. On a certain day the low tide had a height of  $13$  m at  $12.58$  p.m. and the following high tide had a height of  $18$  m at  $6.58$  p.m.

If a ship requires a depth of  $16.5$  m of water before it can leave the harbour, find the latest time on that day that the ship can leave the harbour.

7. (a) Define
- limiting friction,
  - coefficient of friction,
  - angle of friction.

- (b) A uniform rod of length  $2a$  rests on a rough horizontal plane at point  $o$ , and is held inclined at an angle of  $30^\circ$  to the horizontal by a string tied at its top end and to a fixed point  $p$  distant  $2a$  vertically above  $o$ .

If the rod is on the point of slipping, calculate the coefficient of friction between the rod and the plane.



8. (a) Prove that the moment of inertia of a uniform square lamina, of mass  $m$  and side  $2a$ , about an axis through its centre parallel to one of its sides is  $\frac{1}{3}ma^2$ .
- (b) A square lamina  $p, q, r, s$  can turn freely about a horizontal axis through  $p$  perpendicular to the plane of the lamina.
- If the lamina is released from rest when diagonal  $pr$  is horizontal, find its angular velocity when  $pr$  is vertical.
  - What mass must be attached to the lamina at  $r$  so that the combined body will oscillate with period (of small oscillations)

$$2\pi\sqrt{\frac{8a}{3g}}$$

9. (a) State Archimedes' Principle.  
A plastic block of volume  $330 \text{ cm}^3$  has air bubbles in it. The block floats in water with 80% of its volume immersed. If the relative density of the plastic is 1.2, calculate the volume of the air bubbles.
- (b) A cubical block of wood of side 10 cm and of relative density 0.6 floats horizontally in a container of water. Oil of relative density 0.8 is poured on the water until the top of the oil layer is 3 cm below the top of the block.
- How deep is the layer of oil?
  - What is the pressure on the lower face of the block?

10. (a) Solve the differential equation  $x \frac{dy}{dx} = y(1 + y)$   
if  $x = 1$  when  $y = 1$ .

- (b) A particle of mass 8 kg starts from rest and is acted on by a force which increases uniformly in 10s from zero to 16N.
- Prove that  $t$  seconds after the particle begins to move, its acceleration is  $\frac{t}{5} \text{ m/s}^2$ .
  - Prove that, when the particle has moved  $x$  m, its speed is  $v$  m/s, where  $10v^3 = 9x^2$ .