## AN ROINN OIDEACHAIS

## **JUNIOR CERTIFICATE EXAMINATION, 1996**

## MATHEMATICS — HIGHER LEVEL — PAPER 1 (300 marks)

## THURSDAY, 6 JUNE — MORNING, 9.30 to 12.00

Attempt QUESTION 1 (100 marks) and FOUR other questions (50 marks each).

Marks may be lost if necessary work is not clearly shown. Mathematics Tables may be obtained from the Superintendent.

- (i) A distance of 375 metres is travelled in 25 seconds.
   Find the average speed in kilometres per hour.
  - (ii) By selling a car for IR£8840 a loss of 35% is made on the purchase price.

    Calculate the purchase price.
  - (iii) A sphere with radius of length 3 cm has a volume equal to eight times the volume of a sphere with radius of length r cm. Calculate r.
  - (iv) If  $A = \{1,2,3\}$ ,  $B = \{2,3,4\}$  and  $C = \{3,4,5\}$ , find the set

- (v) If  $f: x \to \frac{x}{3} 2$ , what is  $f^{-1}(-5)$ ?
- (vi) Solve for x:

$$3(x-1)^2 - 2(x-1) - 1 = 0$$

- (vii) If  $\frac{1}{b} = \frac{3}{p} \frac{4}{a}$ , express p in terms of a and b.
- (viii) If x \* y = x(3y 5), calculate the value of 3 \* (5 \* 2).
  - (ix) If  $27^{-\frac{1}{3}} = x$ , find the value of x.
  - (x) Using the graph of a quadratic function, or otherwise, find the solution set of the inequality

$$6 - x - x^2 \ge 0, \quad x \in \mathbf{R}.$$


(a) (i) A circle with centre o and radius of length 6 cm contains the points a and b.

Find the area of the shaded region if  $|\angle aob| = 90^{\circ}$ .

Give your answer in terms of  $\pi$ .

(ii) Calculate the total length of the perimeter of the shaded region.

(Take  $\pi = 3.14$ ).



- (b) Wax in the shape of a cylinder with radius of length 4 cm and height 36 cm is melted down. The resulting wax is formed into cone shaped candles. Each candle has height 6 cm and base of radius length 2 cm.
  - (i) Calculate the number of candles that can be made, assuming that no wax is lost.
  - (ii) The candles are placed, base down and in rows of three, in the smallest possible rectangular box. Calculate, in cm<sup>3</sup>, the volume (internal capacity) of the box.
  - (iii) What percentage of the volume of the box is empty? Take  $\pi = 3.14$ . Give your answer correct to the nearest whole number.
- 3. (a) Solve for x:

$$\frac{4}{(x-2)(x+1)} - \frac{2}{x+1} = \frac{3}{x-2}, \quad x \neq 2, \ x \neq -1.$$

(b) Factorise each of the following:

(i) 
$$3x^2 + 2x - 8$$

(ii) 
$$2x^2y - 2xz - 3xy + 3z$$

(iii) 
$$9a^2 - 12ab + 4b^2 - 16c^2$$
.

(c) Solve, correct to 1 place of decimals, the equation

$$5x^2 - 7x - 10 = 0.$$

Hence, or otherwise, find the values of y for which

$$5\left(\frac{1}{y}\right)^2 - 7\left(\frac{1}{y}\right) - 10 = 0$$

and give your answers correct to 1 place of decimals.



The table below gives a summary of the amount of money, in IR£, spent by 90 people at a charity auction:

Amount of money (IR£)	0-10	10-20	20-30	30-50	50-70	70-100
Number of people	26	18	15	12	10	9

(Note: 0-10 means 0 or more but less than 10, etc.)

- (i) Draw a histogram to illustrate the data.(Graph paper is available from the Superintendent).
- (ii) Using mid-interval values, estimate the mean amount spent per person.

  Give your answer correct to the nearest IR£.
- (iii) Copy and complete the cumulative frequency table:

Amount of money (IR£)	<10	<20	<30	<50	<70	<100
Number of people		44				

- (iv) Draw the cumulative frequency curve (ogive) from the table in (iii), putting the number of people on the vertical axis.
- (v) Use the cumulative frequency curve to estimate the interquartile range.
- 5. (a) Graph the function  $f: x \to 9 + 3x 2x^2$  in the domain  $-2 \le x \le 4$ ,  $x \in \mathbb{R}$ . Estimate from your graph
  - (i) the value of f(1.6)
  - (ii) the roots of the equation  $2x^2 3x 10 = 0$
  - (iii) the range of values of x for which

$$7 + 3x - 2x^2 \ge 0.$$

- **(b)**  $f: x \to 3x 1$  and  $g: x \to 3x^2 + 2$  are two functions defined on **R**.
  - (i) Find expressions in x for  $(f \circ g)(x)$  and for  $(g \circ f)(x)$ .
  - (ii) Find the two values of x for which


$$(fog)(x) = (gof)(x).$$




$$3x - 2y = y - 6x$$

$$\frac{5x - 3y + 2}{2} = \frac{x - 2y + 4}{3}$$

- (b) If  $\log_a 2 = p$  and  $\log_a 3 = q$ , express, in terms of p and q,
  - (i)  $\log_a 12$
  - (ii)  $\log_a \frac{9}{4}$



-1 -3



(c) A rectangle of area 72 cm<sup>2</sup> is divided into 4 smaller rectangles. The lengths of the sides are as shown in the diagram.

Write down an equation in x to represent this information.

Solve for x.



