

Fractional integration, differentiation, and weighted Bergman spaces

by

Stephen M. Buckley,¹ Pekka Koskela,¹ and Dragan Vukotić

Abstract: We study the action of fractional differentiation and integration on weighted Bergman spaces and also the Taylor coefficients of functions in certain subclasses of these spaces. We then derive several criteria for the multipliers between such spaces, complementing and extending various recent results. Univalent Bergman functions are also considered.

§0. Introduction

The question of describing the coefficient multipliers between various spaces of analytic functions is an old subject that essentially began with the works of Hardy and Littlewood. Their second paper on fractional integrals [16] in the 1930's contains several results of this kind, even though they do not explicitly mention multipliers. Results involving the integration or differentiation of an analytic function in the unit disc, such as the classical theorem of Littlewood and Paley (see [23] or [3]) which says that if $f \in H^p$ ($p \geq 2$), then f' belongs to the weighted Bergman space A_{p-1}^p , can be viewed as coefficient multiplier results. Some similar results for $p < 2$ were obtained later in [17], [31], and [3]. A related problem also arises in connection with the differentiation operators and Carleson measures (area version): for which positive values of p , q , and for which $\alpha, \beta > -1$, are the quantities

$$\int_{\mathbf{D}} |f(z) - f(0)|^p (1 - |z|^2)^\alpha dA(z) \quad \text{and} \quad \int_{\mathbf{D}} |f'(z)|^q (1 - |z|^2)^\beta dA(z)$$

equivalent? Flett ([14] and [15]) provided the answer in various cases; general questions of this type were also studied recently by Luecking [22]. A complete answer to this question follows from Theorem 1.3 below (which treats arbitrary fractional derivatives).

From the late 50's to the early 70's Flett ([13], [14], [15], etc.) obtained a number of precise results about the order of growth of a function and its (fractional) integrals and derivatives in terms of the Taylor series coefficients of functions using the general spaces

¹ The first author was partially supported by NSF Grant DMS-9207715 and by Forbairt. The second author was partially supported by NSF Grant DMS-9305742 and by the Academy of Finland.
AMS subject classification: Primary 30B10; Secondary 42A45 30D55 30H05
Final version: 19 May 1997

$H(p, q, s)$. Since then, many authors have worked on the theory of multipliers between various spaces of analytic functions. The survey article by Campbell and Leach [9] and a recent paper of Blasco [6] contain many theorems and references. Some other related results can also be found in [1], [4], [11], [12], [19], [21], [25], and [26].

Pavlović [28] found a characterisation of (A^p, A^q) for $0 < p \leq 1, p \leq q < \infty$, in terms of the growth of integral means of a certain function associated with the “candidate” sequence; here, A^p is analytic $L^p(\mathbf{D})$, and (A^p, A^q) is the space of multipliers from A^p to A^q . Since it is not easy to control the growth of a function in terms of its coefficients (see the results in Section 4 on random changes of sign), it is of interest to find necessary or sufficient sequential conditions even in these cases.

In a significant breakthrough, Wojtaszczyk [30] characterised the multipliers (A^p, A^q) , $0 < q \leq 2 \leq p < \infty$, in terms of mixed-norm sequence spaces $l^{\infty, r}$; this result was simplified and generalised by Anderson [2], Jevtić and Jovanović [18], and Blasco [6]. Blasco also described (A^p, A^2) for $0 < p \leq 1$. Vukotić [29] had earlier characterised (A^1, A^2) , and had given several conditions which are either necessary or sufficient for membership of (A^p, A^q) in certain cases, such as $0 < p \leq 2 \leq q < \infty$.

In Section 1, we reduce the study of multipliers between weighted Bergman spaces to that for unweighted ones; this reduction, which we deduce from the work of Flett, appears to have gone unnoticed previously.

One cannot characterise Bergman functions (for $p \neq 2$) in terms of the size of the Taylor coefficients but, in Section 2, we show that this is possible for functions with lacunary or monotonic Taylor sequences (these examples are important for later results).

In Section 3 we discuss the self-multipliers (X, X) for certain spaces X of analytic functions. We show that (X, X) often contains BV , the sequences of bounded variation. Even when this is false, we can almost always say that (X, X) contains the “Taylor form” sequences (which form a subclass of BV); see Section 3 for precise statements.

A careful inspection of the literature reveals a need to justify the free interchange between two forms of fractional integrals and derivatives (those expressed in terms of the special function B and those given by powers of n). This was done in certain cases by Flett [15]; Taylor form multipliers allow us to do so for many function spaces.

In Section 4, we give various (but separate) necessary and sufficient conditions for a sequence to be a multiplier between weighted Bergman spaces. Finally, in Section 5 we briefly consider the action of fractional differentiation on univalent A^p functions; in particular, we generalise part of a well-known result of Brennan [7].

§1. Preliminaries. Fractional Derivatives and Integrals

Unless otherwise stated, analytic functions are assumed to be analytic on the unit disc $\mathbf{D} \subset \mathbf{C}$, and we identify such functions with their sequence of Taylor coefficients at the origin. Writing dA for Lebesgue measure on \mathbf{D} , we are mainly interested in the weighted Bergman spaces A_α^p ($0 < p < \infty, -1 < \alpha$) of analytic functions f for which

$$\|f\|_{A_\alpha^p} \equiv \left(\int_{\mathbf{D}} |f(z)|^p (1 - |z|)^\alpha dA(z) \right)^{1/p} < \infty.$$

We write $A^p = A_0^p$. We shall also use the more general spaces $H(p, q, s)$. Specifically if $0 < p \leq \infty$, $0 < q, s < \infty$, this is defined as the space of analytic functions for which

$$\|f\|_{H(p,q,s)} \equiv \left(\int_0^1 M_p(r, f)^q (1-r)^{sq-1} dr \right)^{1/q} < \infty,$$

where $M_p(r, f)$ is the $L^p(\{z : |z| = r\})$ “norm” of f . We also define $H(p, \infty, s) \equiv H_s^p$ for all $0 < p \leq \infty$, $0 \leq s$, to be the space of analytic functions for which

$$\|f\|_{H_s^p} \equiv \sup_{0 < r < 1} (1-r)^s M_p(r, f) < \infty.$$

It is clear that $A_\alpha^p = H(p, p, (1+\alpha)/p)$ and, in particular, $A^p = H(p, p, 1/p)$. By an “allowable” triple (p, q, s) , we shall simply mean a triple for which $H(p, q, s)$ has been defined above. To avoid cluttered lists of assumptions, we implicitly assume that all triples are allowable when discussing $H(p, q, s)$ spaces; similarly we implicitly assume that $0 < p < \infty$ and $-1 < \alpha < \infty$ for A_α^p spaces.

We say $\lambda = (\lambda_n)_{n=0}^\infty$ is a *multiplier* between the sequence spaces X and Y , written $\lambda \in (X, Y)$, if $\lambda(a_n) \equiv (\lambda_n a_n) \in Y$ whenever $(a_n) \in X$. We define the *fractional derivative* multipliers by $D^t = (n^t)$ for all $t \in \mathbf{R}$. Although notationally unnecessary, we also define the *fractional integral* multipliers by $I^t = D^{-t}$. Here and later, 0^t is taken to mean 0 (even if t is negative); this abuse of arithmetic is harmless, because membership of any space considered below is always an asymptotic property.

Given any sequence $a = (a_k)$, and any $0 < p \leq \infty$, we define the sequence $s(a, p) = (s_n(a, p))_{n=0}^\infty$ by $s_n(a, p) = \left(\sum_{k=2^n}^{2^{n+1}-1} |a_k|^p \right)^{1/p}$ (with the usual alteration for $p = \infty$). If also $0 < q \leq \infty$, the space $l(p, q)$ is defined as the set of all sequences a for which $s(a, p) \in l^q$. Clearly $l(p_1, q_1) \subseteq l(p_2, q_2)$ if $p_1 \leq p_2$, $q_1 \leq q_2$, and also $l(p, p) = l^p$ for all $p > 0$. In general the $l(p, q)$ condition can be viewed as a slight perturbation of the l^p condition (for instance, if $q < \infty$, $(n^{-t}) \in l(p, q)$ if and only if $tp > 1$).

If X is a sequence space and, for every sequence (a_n) , $\lambda(a_n) \in X$ if and only if $\lambda'(a_n) \in X$, we say that λ and λ' are *equivalent on X* . We simply say that λ and λ' are *equivalent* if they are equivalent on all weighted Bergman spaces; obviously, equivalent sequences are multipliers between the same pairs of Bergman spaces. If A and B are any positive quantities, we write $A \approx B$ if $A/C \leq B \leq CA$, where C is any positive constant whose exact value does not matter.

We first gather together some known results for the spaces $H(p, q, s)$ and A_α^p in a pair of lemmas. Parts (i)–(iii) of our first lemma (see [6]) are fairly elementary; (iv) is deeper and was proven by Flett ([14] for $s = 0, q = \infty$ and [15, Theorem 6] for all other cases). Part (v) follows easily by rewriting Theorem 5(iii) of [15]. The second lemma is taken from [27] (parts (i)–(iv), (vi)), while part (v) is from [17].

Lemma 1.1. *If $p, q, s, s+t > 0$, $p_1 \leq p_2$, $q_1 \leq q_2$, and $0 < s_1 \leq s_2$, then*

- (i) $H^{p_2} \subset H(p_2, q_1, s_1) \subset H(p_1, q_2, s_2) \subset H_{s_2}^{p_1}$.
- (ii) $H(p_1, q, s) \subset H(p_2, q, s + p_1^{-1} - p_2^{-1})$.
- (iii) $H(p, q_2, s) \subset H(p, q_1, s + q_1^{-1} - q_2^{-1})$.
- (iv) $D^t \in (H(p, q, s), H(p, q, s+t))$.
- (v) $I^s \in (H(p, p, s), H^p)$, for $0 < p \leq 2$.

Lemma 1.2. Let $f(z) = \sum_{n=0}^{\infty} a_n z^n$.

- (i) If $0 < p \leq 1$ and $f(z) \in A_{\alpha}^p$, then $a_n = o(n^{\frac{\alpha+2}{p}-1})$.
- (ii) If $1 \leq p < \infty$ and $f(z) \in A_{\alpha}^p$, then $a_n = o(n^{\frac{\alpha+1}{p}})$.
- (iii) If $f(z) \in A_{\alpha}^p$, then $\sum_{n=1}^{\infty} n^{p-\alpha-3} |a_n|^p < \infty$ whenever $0 < p \leq 2$, and the converse holds whenever $p \geq 2$.
- (iv) If $f(z) \in A_{\alpha}^p$, then $\sum_{n=1}^{\infty} n^{-\alpha-1} |a_n|^p < \infty$ whenever $p \geq 2$, and the converse holds whenever $0 < p \leq 2$.
- (v) Suppose p, p' are conjugate indices. If $f(z) \in A^p$, then $\sum_{n=1}^{\infty} n^{1-p'} |a_n|^{p'} < \infty$ whenever $1 < p \leq 2$, and the converse holds whenever $p \geq 2$.
- (vi) If $q < p$, then $A_{\alpha}^p \subset A_{\beta}^q$ for all β satisfying $(\beta + 1)q^{-1} > (\alpha + 1)p^{-1}$.

We now give an answer to the question on equivalence of two integrals mentioned in the introduction. The next couple of results follow easily from Lemmas 1.1 and 1.2, but do not seem to have been noticed or written down before in full generality. In particular, they generalise earlier results of Flett [15]. We would like to stress that parts (c) and (d) of Corollary 1.4 completely reduce the study of multipliers and coefficient estimates on weighted Bergman spaces to the unweighted case. All indices and inequalities in these first two results are sharp, and one cannot replace the “ $<$ ” by “ \leq ” in parts (b) of 1.3 and 1.4, but we postpone justifications of sharpness until later (see comments after Proposition 2.1 and Lemma 3.8).

Theorem 1.3. If $-1 < \alpha, \beta < \infty$, $t \in \mathbf{R}$, and $0 < p, q \leq \infty$, then $D^t \in (A_{\alpha}^p, A_{\beta}^q)$

- (a) if $p \leq q$ and $(2 + \beta)q^{-1} \geq (2 + \alpha)p^{-1} + t$ or,
- (b) if $p > q$ and $(1 + \beta)q^{-1} > (1 + \alpha)p^{-1} + t$.

Proof. The special case $p = q$ of Lemma 1.1(iv) gives us $D^t \in (A_{\alpha}^p, A_{\alpha+tp}^p)$ for all $\alpha > -1$, $t \in \mathbf{R}$, $p > 0$. But by Lemma 1.1(ii),

$$A_{\alpha+tp}^p = H(p, p, (1 + \alpha)p^{-1} + t) \subset H(q, q, (2 + \alpha)p^{-1} - q^{-1} + t) = A_{\beta}^q$$

for any $q \geq p$, giving us (a). Similarly, (b) follows by using Lemmas 1.1(iv) and 1.2(vi). \square

Corollary 1.4.

- (a) If $p \leq q$, and $t \geq (2 + \alpha)(p^{-1} - q^{-1})$, then $I^t \in (A_{\alpha}^p, A_{\alpha}^q)$.
- (b) If $p > q$, and $t < (1 + \alpha)(q^{-1} - p^{-1})$, then $D^t \in (A_{\alpha}^p, A_{\alpha}^q)$.
- (c) $(a_n) \in A_{\alpha}^p$ if and only if $(n^{-\alpha/p} a_n) \in A^p$.
- (d) $(\lambda_n) \in (A_{\alpha}^p, A_{\beta}^q)$ if and only if $(n^{\alpha p^{-1} - \beta q^{-1}} \lambda_n) \in (A^p, A^q)$.

Proof. (a) and (b) are special cases of the corresponding parts of Theorem 1.3; (c) follows immediately from (a), and (d) from (c). \square

Since the indices in Theorem 1.3 are best possible (as we shall see), the case $t = 0$ gives the best possible imbeddings of the form $A_{\alpha}^p \subset A_{\beta}^q$. With one exception, these are as given in Theorem 1 of [27]. The exception is their imbedding 3° (and the equivalent 7°) which says that if $(1 + \alpha)/p > (1 + \beta)/q$ and $(2 + \alpha)/p < (2 + \beta)/q$ then $A_{\alpha}^p \subset A_{\beta+1}^q$.

In this case, we must have $p > q$ and so Theorem 1.3(b) tells us that $A_\alpha^p \subset A_\gamma^q$ for all $\gamma > \gamma_0 = (1 + \alpha)qp^{-1} - 1$. Note that $\beta < \gamma_0 < \beta + 1$ in all cases.

Suppose f, g have Taylor coefficients $(a_n), (b_n)$ respectively. We shall make use of the pairing

$$\langle f, g \rangle = \frac{1}{\pi} \int_{\mathbf{D}} f \bar{g} dA = \sum_{n=0}^{\infty} \frac{a_n \bar{b}_n}{n+1}.$$

Suppose $1 < p < \infty$. It is well-known (e.g., see Theorem 1.16 of [5]) that the dual $(A^p)^*$ of A^p with respect to this pairing is $A^{p'}$ where $p' = p/(p-1)$ is the index dual to p ; furthermore the $(A^p)^*$ -norm of the linear functional corresponding to $g \in A^{p'}$ is comparable with $\|g\|_{A^{p'}}$. If also $1 < q < \infty$ and $q' = q/(q-1)$, it follows that $\lambda \in (A^p, A^q)$ if and only if $\bar{\lambda} \in (A^{p'}, A^{q'})$. But it is clear that $(a_n) \in A^r$ if and only if $(\bar{a}_n) \in A^r$ ($0 < r < \infty$), and so we deduce that $(A^p, A^q) = (A^{q'}, A^{p'})$, a useful duality result that we shall use later in this paper. It also follows by duality (although now the pairing $\sum_{n=0}^{\infty} a_n \bar{b}_n$ is more natural) that $(H^p, H^q) = (H^{q'}, H^{p'})$.

§2. Functions with lacunary or monotonic Taylor coefficients

We shall later look at necessary and sufficient conditions for multipliers between Bergman spaces. For this analysis, it will be useful to build up a stock of examples and non-examples of functions in A_α^p .

Our first class of examples are *lacunary sequences* (recall that we are identifying analytic functions with their sequences of Taylor coefficients). We define the k th dyadic block B_k to be the set of integers from 2^{k-1} to $2^k - 1$ inclusive, if $k > 0$, and $B_0 = \{0\}$. Also let \tilde{B}_k be the larger set of integers from $2^{k-2} + 1$ to $2^k - 1$ inclusive, if $k > 1$, and $\tilde{B}_0 = \{0\}$, $\tilde{B}_1 = \{0, 1\}$. Our lacunary sequences include all the usual lacunary (a_n) , where $a_n = 0$ if n is not a power of 2 but, in view of our later needs, we define a *lacunary sequence* to be any sequence which has at most one non-zero term with index in any one block B_k (and hence at most two non-zero terms corresponding to each \tilde{B}_k).

The following useful characterisation of which lacunary sequences lie in A_α^p may be known, but we include a proof for lack of a specific reference.

Proposition 2.1. *Suppose (a_n) is a lacunary sequence. Then for any $0 < p < \infty$, $(a_n) \in A_\alpha^p$ if and only if $\sum_{n=1}^{\infty} n^{-\alpha-1} |a_n|^p < \infty$.*

Proof. The main idea is to write down the norm of a function as the sum of norms over the dyadic blocks, in the spirit of what is often called the Littlewood-Paley theory.

We may as well assume $\alpha = 0$ by Corollary 1.4(c). As a special case of [6, (2.3) and Theorem 3.1], we have

$$\|f\|_{A^p} \approx \left[\sum_{k=0}^{\infty} 2^{-k} \|w_k * f\|_{H^p}^p \right]^{1/p}, \quad (2.2)$$

where, for each k , w_k is a sequence whose terms are non-negative and bounded by 1, and whose terms are zero for indices outside \tilde{B}_k ; additionally the sum over k of the n th terms

of the w_k 's is 1 for all n . If $p > 1$, one can simply choose $w_k = \Delta_k$, the characteristic multiplier which is 1 on B_k and zero elsewhere. For $p \leq 1$, it is shown in [25] that this simple but rough cutoff still gives one half of (2.2) (the right-hand side dominates the left) but to get the full strength result, one must choose a smoother cutoff for large k .

The proposition therefore follows easily from the following two inequalities:

$$\|w_k * f\|_{H^p}^p \leq C_1 \sum_{n \in \tilde{B}_k} |a_n|^p, \quad (2.3a)$$

$$\sum_{j=k}^{k+1} \|w_j * f\|_{H^p}^p \geq C_2 \sum_{n \in B_k} |a_n|^p, \quad (2.3b)$$

where C_1 and C_2 are independent of k . Since there are at most two non-zero terms with index in \tilde{B}_k , (2.3a) is easily seen to be valid with $C_1 = 2$. (2.3b) is almost as easy to verify, so we leave it as an exercise with two hints: (1) if $n \in B_k$ then the n th coefficient of either w_k or w_{k+1} is at least $1/2$; (2) the H^p norm of a power series remains unchanged if we divide by a power of z to make the constant term non-zero. \square

Suppose (λ_n) is lacunary and all of its terms are 0 or 1. It is now obvious that $(\lambda_n n^{1/p})$ is in A^q if and only if $0 < q < p$. Also $(\lambda_n n^{1/p} \log^{-t} n) \in A^p$ if and only if $tp > 1$. Considering the action of fractional differentiation on this last example shows that one cannot replace the inequality in Theorem 1.3(b) or Corollary 1.4(b) by equality.

As another application of Theorem 1.3, one can prove that Lemma 1.2(ii) is best possible in the sense of Duren-Taylor: if δ_n is any sequence tending to zero, $p \geq 1$, and $\alpha > -1$, then there are sequences $(a_n) \in A_\alpha^p$ such that $a_n \neq O(\delta_n n^{\frac{\alpha+1}{p}})$. This was shown in [27] when $1 \leq p < 2$ using lacunary sequences and the converse part of Lemma 1.2(iv), but the argument works also for $p > 2$ since Proposition 2.1 says that the converse part of 2.1 is also valid for $p > 2$ if (a_n) is lacunary (also the example given in [27] to prove that the exponent in 1.2(ii) is best possible when $p > 2$ does not work as claimed).

The second important class of examples we wish to examine are monotone sequences. For $p > 1$, we characterise the monotonic sequences which are in A^p .

Proposition 2.4. *Suppose $a_n \geq 0$, $1 < p, C < \infty$ and that either*

- (i) (a_n) is monotonic, or
- (ii) $(a_n)_{n \in B_k}$ is monotonic and, for all $n, m \in B_k$, $a_m \leq C a_n$.

Then $(a_n) \in A_\alpha^p$ if and only if $\sum_{n=1}^{\infty} n^{p-3-\alpha} |a_n|^p < \infty$.

Proof. As usual we may assume $\alpha = 0$. We claim that for all $n \in B_k$, $\|\sum_{n \leq m \in B_k} z^m\|_{H^p} \leq C 2^{k(p-1)/p}$. One approach to justifying this is to write the sum as $(z^{2^k} - z^n)/(z - 1)$ and find upper bounds for this expression for $z = e^{i\theta}$ in the ranges $|\theta| \leq 2^{-k}$ and $2^{-j-1} < |\theta| \leq 2^{-j}$, $j = 0, \dots, k$; in any case, we leave the details to the reader. Using monotonicity and the summation by parts formula, we have

$$\sum_{n \in B_k} a_n z^n = a_{2^{k-1}} \sum_{n \in B_k} z^n + \sum_{2^{k-1} < n \in B_k} (a_n - a_{n-1}) \sum_{n \leq m \in B_k} z^m,$$

and therefore $\|\sum_{n \in B_k} a_n z^n\|_{H^p} \leq C 2^{k(p-1)/p} (a_{2^{k-1}}^p + a_{2^k-1}^p)^{1/p}$.

As previously mentioned, $\|f\|_{A^p} \leq C [\sum_{k=0}^{\infty} 2^{-k} \|\Delta_k * f\|_{H^p}^p]^{1/p}$ for all $p > 0$, where Δ_k is the characteristic multiplier of B_k . Thus

$$\|f\|_{A^p}^p \leq C \left[\sum_{k=0}^{\infty} 2^{k(p-2)} (a_{2^{k-1}}^p + a_{2^k-1}^p) \right],$$

and one half of the equivalence follows.

Conversely, suppose $f \in A^p$. By Lemma 1.2(iii), it suffices to prove the desired implication for $p > 2$. Using monotonicity and one direction of (2.2), the proof follows easily from the estimate $\|\sum_{n \in B_k} a_n z^n\|_{H^p} \geq C 2^{k(p-1)/p} \min_{n \in B_k} a_n$. This last inequality is immediate if one considers a lower bound on the real part of the left-hand sum on the arc $\{z = e^{i\theta} : |\theta| < 2^{-k}\}$. \square

§3. Self-multipliers

A *self-multiplier* of X is a member of (X, X) . Using Stirling's formula, we see that

$$A \sum_{n=1}^{\infty} |a_n|^2 / n^{1+\alpha} \leq \|f\|_{A_\alpha^2}^2 \leq B \sum_{n=1}^{\infty} |a_n|^2 / n^{1+\alpha},$$

for some $0 < A \leq B < \infty$ ($A = B = 1$ when $\alpha = 0$). Consequently, the self-multipliers of A_α^2 are precisely all bounded sequences. By Corollary 1.4(d), we always have $(A_\alpha^p, A_\alpha^p) = (A^p, A^p)$, but no simple description of (A^p, A^p) is known when $p \neq 2$. In this section, we discuss certain useful types of examples and non-examples of such self-multipliers.

We recall that a sequence space X is said to be *solid* if $(b_n) \in X$ whenever $|b_n| \leq |a_n|$, $(a_n) \in X$. The spaces $l^{p,q}$ and c_0 are examples of solid spaces. We shall see later that (A^p, A^p) is not solid when $p \neq 2$, and so it cannot be expressed in terms of any of the standard (solid) sequence spaces.

Suppose $X^p = H^p$ or A_α^p , $0 < p < \infty$, $-1 < \alpha < \infty$. By considering vectors of the form z^n , the Closed Graph Theorem immediately implies that $(X^p, X^p) \subset l^\infty$ (and containment is strict whenever $p \neq 2$; see Corollary 4.4).

The shift operators σ^- and σ^+ , as usual, are defined by

$$\begin{aligned} \sigma^-(a_0, a_1, a_2, \dots) &= (a_1, a_2, \dots), \\ \sigma^+(a_0, a_1, a_2, \dots) &= (0, a_0, a_1, a_2, \dots). \end{aligned}$$

We call a topological vector space of analytic functions an *M-space* if it is metrisable by an invariant metric d "controlled by" $\{M_p(r, f) : 1/2 < r < 1\}$ for some $p > 0$, in the sense that if $M_p(r, f) \leq 2M_p(r, g)$ for all $1/2 < r < 1$ then $d(f, 0) \leq Cd(g, 0)$. For example, $H(p, q, s)$ is an M-space for any allowable (p, q, s) . We record here a simple lemma which we shall use implicitly wherever we need it.

Lemma 3.1. σ^+ and σ^- are bounded on all M -spaces.

Proof. Since $\sigma^+(f(z)) = zf(z)$, we have $M_p(r, \sigma^+(f)) \leq M_p(r, f)$ for all $0 < r < 1$, and so σ^+ is bounded. Similarly, $\sigma^-(f(z)) = (f(z) - f(0))/z$, and $M_p(r, f - f(0)) \leq 2M_p(r, f)$ (since $f(0) \leq M_p(r, f)$ for all $p, r > 0$). Thus $M_p(r, \sigma^-(f)) \leq 4M_p(r, f)$ for any $1/2 < r < 1$, and so σ^- is bounded. \square

If X and Y are M -spaces, then $\lambda = (\lambda_n)_{n=0}^\infty \in (X, Y)$ if and only if $(\lambda_{n+1})_{n=0}^\infty \in (X, Y)$ (since the latter multiplier is simply $\sigma^- \lambda \sigma^+$). We shall use Lemma 3.1 mainly in this form, as a convenience to simplify later statements. For example, $((n+1)^t)$ and (n^t) are equivalent for all $t \in \mathbf{R}$.

We say that a vector space of analytic functions X has the *small multiplier property* if there exists $r > 0$ such that if $\lambda_n = O(n^{-r})$, then $\lambda = (\lambda_n) \in (X, X)$. We say that a sequence (λ_n) has *Taylor form* if there are sequences $(a_i), (C_i)$ such that,

$$\forall k, n > 0: \quad \left| \lambda_n - \sum_{i=0}^k a_i (n+1)^{-i} \right| \leq C_k (n+1)^{-k-1}.$$

We refer to the sequence $(p_{k,n})_{n=1}^\infty \equiv (\sum_{i=1}^k a_i (n+1)^{-i})$ as the k th *Taylor polynomial* of λ , $(\lambda_n - p_{k,n})_{n=1}^\infty$ as the k th *remainder*, and $(a_i (n+1)^{-i})_{n=1}^\infty$ as the i th *Taylor summand*. Taylor form is a purely asymptotic property: it is invariant under changes of a finite number of terms. Also, the above type of expansion in negative powers of $n+1$ is equivalent to the corresponding expansion in negative powers of n but is more convenient for the next lemma.

The following lemma is simple, but very useful—it is used (e.g., in Lemma 3.8, and also to study compact multiplier operators in [8]) to prove that certain pairs of multipliers are equivalent on most function spaces when intuitively we expect this to be true.

Lemma 3.2. *Suppose that X is a vector space of analytic functions which has the small multiplier property, and that $((n+1)^{-1}) \in (X, X)$. Then all multipliers which have Taylor form are self-multipliers on X .*

Proof. Let λ have Taylor form. The k th Taylor summand of λ is a self-multiplier of X for $k = 0$ since X is a vector space, and for all $k > 0$ by iterating the fact that $((n+1)^{-1}) \in (X, X)$. By the small multiplier property, the k th remainder of λ is a self-multiplier of X if k is sufficiently large. It follows that $\lambda \in (X, X)$. \square

The hypotheses for X in the above lemma are quite weak. For example, we have the following lemma.

Lemma 3.3. *All $H(p, q, s)$ spaces satisfy the hypotheses of Lemma 3.2.*

Proof. To see that $X \equiv H(p, q, s)$ has the small multiplier property, let us show that $\lambda \in (X, H^\infty)$ whenever λ_n decreases at a sufficiently fast rate. By Lemma 1.1, $H^p \subset H(p, q, t) \subset H(2, 2, s)$ where $s = t + \max(0, 1/p - 1/2) + \max(0, 1/2 - 1/q)$, and so it suffices to prove the result for A_α^2 , for all $\alpha > -1$. If $(a_n) \in A_\alpha^2$ then $a_n = O(n^{(1+\alpha)/2})$, and so $\lambda \in (A_\alpha^2, H^\infty)$ if $\lambda_n = O(n^{-2-\alpha/2})$, say.

The fact that $((n+1)^{-1}) \in (X, X)$ follows from Lemma 1.1(iv) in all cases except $X = H^p$. In this case, it is true by a theorem of Hardy and Littlewood [10, Theorem 5.12]. \square

Let $B(\cdot, \cdot)$ be the classical Beta function. The alternative fractional integration and differentiation multipliers $\tilde{I}^t = ((B(n+1, t))$ and $\tilde{D}^t = (1/B(n+1, t))$, are used by some authors. We shall use Lemma 3.2 to prove that the two types of fractional operations are equivalent on $H(p, q, s)$. This equivalence was proved in certain cases by Flett [15, Theorem 7]. Here we show that if X is any space of analytic functions satisfying the hypotheses of Lemma 3.2 then these operators are equivalent on X .

Lemma 3.4. *If $\lambda \equiv (\lambda_n) = (n^t B(n+1, t))_{n=1}^\infty$ for some $t > 0$, then both λ and $\lambda^{-1} \equiv (1/\lambda_n)$ have Taylor form.*

Proof. We recall the following asymptotic expansion of $\Gamma(x+1)$ (see, for instance, [24, p.380]):

$$\Gamma(x+1) = \sqrt{2\pi} x^{x+1/2} e^{-x} \left(1 + \frac{1}{12x} + \dots + \frac{a_k}{x^k} + O(x^{-k-1})\right).$$

Thus

$$\frac{n^t B(n+1, t)}{\Gamma(t)} = e^t \cdot \left(\frac{n}{n+t}\right)^{n+t+1/2} \cdot \frac{1 + \frac{1}{12n} + \dots + \frac{a_k}{n^k} + O(n^{-k-1})}{1 + \frac{1}{12(n+t)} + \dots + \frac{a_k}{(n+t)^k} + O(n^{-k-1})}.$$

Thus, to show that λ has Taylor form, it suffices to show that the middle factor has Taylor form. Since

$$x \mapsto (1+tx)^{-t-1/2-1/x} = e^{-\log(1+tx)(t+1/2+1/x)}$$

has a Taylor series expansion about zero (with radius of convergence $1/t$), the required result follows by letting $x = 1/n$. Finally, it is clear from the above that λ^{-1} also has Taylor form. \square

Corollary 3.5. *Suppose X is a vector space of analytic functions satisfying the assumptions of Lemma 3.2, and that f is an analytic function. Then for any $t > 0$, $I^t f \in X$ if and only if $\tilde{I}^t f \in X$, and $D^t f \in X$ if and only if $\tilde{D}^t f \in X$.*

We denote by BV the classical space of (complex) *sequences of bounded variation*:

$$BV = \{(\lambda_n)_{n=0}^\infty : \|(\lambda_n)\|_{BV} = |\lambda_0| + \sum_{n=0}^\infty |\lambda_{n+1} - \lambda_n| < \infty\}.$$

As is also well-known, a sequence belongs to BV if and only if both its real and imaginary parts can be written as differences of two bounded monotonic sequences. Note that BV is not solid and that all Taylor form multipliers have bounded variation (but there are many BV sequences, such as $((-1)^n/(n+1)^2)$, which do not have Taylor form).

Unlike Taylor form multipliers, BV sequences are not necessarily self-multipliers of A^p . In fact, we shall see that $BV \subset (A^p, A^p)$ if and only if $p > 1$. Let us first prove that $BV \subset (X, X)$ when X is any one of many Banach spaces of analytic functions. For this we need a lemma which is due (at least partially) to Zhu [32].

Lemma 3.6. For $X = A^p$ or H^p , $1 < p < \infty$ and $f \in X$, the sequence $(P_n f)_{n=0}^\infty$ of Taylor polynomials converges to f in the norm of X . The statement is false for $p = 1$.

Proposition 3.7. Let $(X, \|\cdot\|)$ be a Banach space of analytic functions in the unit disc \mathbf{D} such that for every $f \in X$ the sequence $(P_n f)_{n=1}^\infty$ of Taylor polynomials converges to f in the norm of X . Then $BV \subset (X, X)$, and the inclusion is strict if the involution $Tf(z) = f(-z)$ is bounded on X . In particular, BV is a proper subset of (X, X) if $X = A^p$ or H^p , $1 < p < \infty$.

Proof. Let $f(z) = \sum_{k=0}^\infty a_k z^k \in X$, and choose $M > 0$ such that $\|(\lambda_n)\|_{BV} \leq M$. Given $\epsilon > 0$, we have for sufficiently large n that

$$\left\| \sum_{k=n}^\infty a_k z^k \right\| < \frac{\epsilon}{2M}$$

(such a choice is possible by our assumption on $P_n f$). A summation by parts yields

$$\left\| \sum_{k=n}^\infty \lambda_k a_k z^k \right\| = \left\| \sum_{k=n+1}^\infty (\lambda_{k+1} - \lambda_k) \cdot \sum_{j=k}^\infty a_j z^j - \lambda_n \sum_{k=n}^\infty a_k z^k \right\| < M \cdot \frac{\epsilon}{2M} + M \cdot \frac{\epsilon}{2M} = \epsilon,$$

for all sufficiently large n . Thus, $\sum_{k=0}^\infty \lambda_k a_k z^k \in X$, which proves the claim. In the case $X = A^p$ or H^p , $1 < p < \infty$, the statement follows from Lemma 3.6.

The involution T corresponds to the sequence $((-1)^n)$ which is not in BV , and so the inclusion is strict if $T \in (X, X)$. Obviously, T is bounded on all $H(p, q, s)$ spaces, including A^p and H^p , $p > 0$. \square

We now define the one-parameter family $f_t(z) = (1 - z)^{-t}$, for all $t > 0$. Simple calculations show that $f_t \in A^p$ if and only if $tp < 2$, and $f_t \in H^p$ if and only if $tp < 1$. Since the n th coefficient of f_t is $1/nB(n, t)$, our results on Taylor form multipliers tell us that $f_t \in A^p$ if and only if $D^{t-1} \in A^p$ (i.e. if and only if $\sum_{n=1}^\infty n^{t-1} z^n \in A^p$), and similarly for H^p .

Lemma 3.8. $D^t \in A^p$ if and only if $(t+1)p < 2$, and $D^t \in H^p$ if and only if $(t+1)p < 1$.

Proof. The lemma is trivially true if $t < -1$, and the preceding comments prove the case $t > -1$. Finally $D^{-1} = (1 - z)[\log(1 - z) - 1]$, which clearly lies in H^∞ . \square

Using D^s as a counterexample is essentially equivalent to using f_{s+1} , but the use of D^s minimises the need to mention Taylor form multipliers. As a first application, it is now clear that the bounds on t in 1.3(a) and 1.4(a) are best possible.

According to Proposition 3.7 every sequence of bounded variation is in (A^p, A^p) for all $p > 1$. Our next theorem tells us that this is false if $p \leq 1$. The proof gives two different methods of constructing counterexamples: the first is simpler but works only if $p < 1$, while our $p = 1$ method is easily adapted to handle all $p \leq 1$, but is less explicit and involves more background work. The case $p < 2/3$ can also be proved from Theorem 4.5 below, a task we leave to the interested reader.

Theorem 3.9. For all $0 < p \leq 1$, there exist monotonic c_0 sequences not in (A^p, A^p) .

Proof. Assume first that $p < 1$. The sequence (n^s) is in A^p for all $s < 2/p - 1$ but, if we perturb its 2^k -th coefficient for all $k > 1$ by subtracting $2^{k/p}$, the resulting sequence is not in A^p by Proposition 2.1. Consequently, $\lambda = (\lambda_n) \notin (A^p, A^p)$ if

$$\lambda_n = \begin{cases} 1 - 2^{k(1/p-s)}, & n = 2^k, k > 1, \\ 1 & \text{otherwise.} \end{cases}$$

Thus if we choose any s in the interval $(1/p, 2/p - 1)$, the resulting λ is a non-multiplier of bounded variation. A monotonic decreasing c_0 non-multiplier is then easily found. One example is the function $f = (\mu_n)$ where

$$\mu_n = \begin{cases} \sum_{j=1}^{\infty} 2^{j(1/p-s)}, & \text{if } n < 2, \\ \sum_{j=k+1}^{\infty} 2^{j(1/p-s)}, & \text{if } n \in B_k, k > 1. \end{cases}$$

To see this, note that the n th Taylor coefficient of $1 + f(z) - zf(z)$ is λ_n .

This leaves the case $p = 1$. It suffices to find a bounded variation non-multiplier (since such a sequence can be written as a difference of two monotonic c_0 sequences, at least one of which must be a non-multiplier). Suppose that no such sequence exists and so, by the Closed Graph Theorem, all BV sequences are bounded as (multiplier) operators on A^1 .

We shall need a slight variant of the characterisation of A^1 given by the $p = 1$ case of (2.2). As given earlier, the functions w_k have non-zero terms only for indices in the range 2^{k-2} to 2^k . This is because the n th term of w_k , which we denote by $w_{k,n}$ equals $\phi(n/2^{k-2})$, where $\phi(t) = \omega(t/2) - \omega(t)$ and ω is any infinitely differentiable function defined on the interval $[0, \infty)$ satisfying $\omega(t) = 1$ if $t \leq 1$, $\omega(t) = 0$ if $t \geq 2$, and $0 \leq \omega(t) \leq 1$ if $1 \leq t \leq 2$. If we widen the allowable class of functions to include all smooth ω which are 1 for $t \leq 1$, 0 for $t \geq 4$, and take on intermediate values on the interval $[1, 4]$, (2.2) is still valid since the proof in [19] of the crucial inequality $\|w_k * f\|_{H^1} \leq \|f\|_{H^1}$, which is needed in [6] to prove (2.2), goes through essentially unchanged (note that our w_k corresponds to w_{k-1} in the other two papers).

Suppose therefore that we choose such an ω so that it is strictly increasing on the interval $[1, 4]$. It follows that the resulting function ϕ has a positive minimum on the interval $[1, 2]$ and hence that there exists $0 < c < 1$ such that for all $k > 1$ and all $n \in B_k$, the n th term of w_k lies between c and 1. Also, it is easy to show (see [25] for example) that if the Taylor coefficients of f are zero outside the range from m to n then $M_p(r, f)/\|f\|_{H^p}$ lies between r^n and r^m ; hence it is true in general that $\|w_k * f\|_{H^1} \approx 2^k \|w_k * f\|_{A^1}$.

By (2.2), we have $\|w_k * f\|_{H^1} \leq C2^k \|f\|_{A^1}$ (for some constant C independent of k). On the other hand, functions f_k are given in [25] for all $k > 0$ such that all their Taylor coefficients with index in B_k are positive, and $\|\Delta_k f_k\|_{H^1} \geq Ck2^k \|f\|_{A^1}$. Letting $\lambda_k = (\lambda_{k,j})_{j=0}^{\infty}$ be defined by $\lambda_{k,j} = 1/w_{k,j}$, if $j \in B_k$, and $\lambda_{k,j} = 0$ otherwise, we see that $\lambda_k(g_k) = \Delta_k f_k$, where we define $g_k = w_k * f_k$. By construction, $\|\lambda_k\|_{BV} \leq T + 2/c$, where T is the total variation of the function $1/\phi$ on $[1, 2]$; in particular, the total variation of λ_k has a bound independent of k . Writing $\lambda = \sum_{k=1}^{\infty} k^{-2} \lambda_{k,3}$, it follows that $\lambda \in BV$, but $\|\lambda(g_{k,3})\|_{A^1} \geq Ck \|g_{k,3}\|_{A^1}$, giving the required contradiction. \square

Our BV -related results can be summarised as follows.

Theorem 3.10. *Each of the following statements about A^p is true if and only if $p > 1$.*

- (a) $BV \subset (A^p, A^p)$.
- (b) Every monotone (real) c_0 sequence is a self-multiplier of A^p .
- (c) If $f \in A^p$, the sequence $(P_n f)_{n=0}^\infty$ of Taylor polynomials is norm-convergent to f .

§4. Sequential estimates for Bergman space multipliers

In this section, we investigate (A_α^p, A_β^q) . The first statement gives a complete characterisation in the case $0 < q \leq 2 \leq p < \infty$. The unweighted version ($\alpha = \beta = 0$) of this is due to Wojtaszczyk [30]. The weighted version given here is due to Anderson [2], who gave a considerably simpler and more elementary proof (of course in light of Corollary 1.4, the unweighted result immediately gives the weighted result). Elementary proofs in general situations were also given in [18] and [6]. Here and subsequently, 0^{-1} is defined to be ∞ .

Lemma 4.1. *Suppose $0 < q \leq 2 \leq p < \infty$, and $\alpha, \beta > -1$. Then $\lambda = (\lambda_n) \in (A_\alpha^p, A_\beta^q)$ if and only if $(n^{-r} \lambda_n) \in l(\infty, s^{-1})$, where $s = q^{-1} - p^{-1}$ and $r = (1 + \beta)q^{-1} - (1 + \alpha)p^{-1}$.*

Lemma 4.1 tells us that (A^p, A^q) is solid if $p \geq 2 \geq q$. The next two theorems, which deal with fractional differentiation composed with a random change of sign, will imply that this is false (for Hardy as well as Bergman spaces) if p, q are either both less than 2 or both greater than 2.

Theorem 4.2. *Suppose that either $0 < p, q \leq 2$, $p \neq 2$, or $2 \leq p, q < \infty$, $q \neq 2$. In the first case, we define $t_0 = 1/q + 1/2 - 2/p$, and in the second case we define $t_0 = 2/q - 1/2 - 1/p$. If $t > t_0$, then for almost every choice of signs $\epsilon_n = \pm 1$, $(\epsilon_n n^t) \notin (A^p, A^q)$.*

Proof. Consider first the case $p < 2$, $q \leq 2$. Let $0 < \delta \leq 1/p - 1/2$ be fixed but otherwise arbitrary. Lemma 3.8 implies that $D^{2/p-1-\delta} \in A^p \setminus A^2$. By Lemma 1.1(v) and the upper bound on δ , we deduce that $D^{1/p-1-\delta} \in H^p \setminus H^2$.

If $(\epsilon_n n^t) \in (A^p, A^q)$ for $t = 1/q + 1/2 - 2/p + \delta$, one immediately obtains $(\epsilon_n n^{1/q-1/2}) \in A^q$. Using Lemma 1.1(v) again, we see that $(\epsilon_n n^{-1/2}) \in H^q$. Since $(n^{-1/2}) \notin A^2$, a theorem of Littlewood [10, Theorem A5] tells us that for almost every choice of signs $\epsilon_n = \pm 1$, $(\epsilon_n n^{-1/2})$ has radial limit almost nowhere (and hence cannot lie in H^s for any $s > 0$).

Since $(A^p, A^q) = (A^{q'}, A^{p'})$ where p', q' are conjugate to p, q , the case $p \geq 2, q > 2$ follows by duality. \square

Theorem 4.3. *Suppose that $0 < p, q < \infty$. Let $t_0 = \min\{1/2 - 1/p, 1/q - 1/2, 0\}$. If $t > t_0$, then for almost every choice of signs ϵ_n , $(\epsilon_n n^t) \notin (H^p, H^q)$.*

Proof. Fix $\delta > 0$. By Lemma 3.8, $D^{1/p-1-\delta} \in H^p$, and so it follows as in Theorem 4.2 that $(\epsilon_n n^{\delta+1/2-1/p}) \notin (H^p, H^q)$ for almost every (ϵ_n) . If $p, q > 1$, then by duality, $(\epsilon_n n^{\delta+1/q-1/2}) \notin (H^p, H^q)$ for almost every (ϵ_n) (and note that $1/q - 1/2 < \min\{1/2 - 1/p, 0\}$ forces $p, q > 1$). Finally, to see that $(\epsilon_n n^\delta) \notin (H^p, H^q)$, we note that $(n^{-1/2-\delta/2}) \in H^2$, and so $(\epsilon'_n n^{-1/2-\delta/2}) \in H^p$ for some fixed choice of signs (ϵ'_n) , but $(n^{-1/2+\delta/2}) \notin H^2$, and so $(\epsilon_n \epsilon'_n n^{-1/2+\delta/2}) \notin H^p$ for almost all (ϵ_n) . \square

Corollary 4.4. *If $0 < p, q < 2$ or $2 < p, q < \infty$, then (A^p, A^q) and (H^p, H^q) are non-solid.*

Proof. It is an easy consequence of Theorem 1.3 that $D^t \in (A^p, A^q)$ for all $t > t_0$ whenever p, q, t_0 are as in Theorem 4.2. Thus, (A^p, A^q) is not solid if p, q are as hypothesised.

Letting $t_1 = 0$ if $q \leq p$, and $t_1 = 1/q - 1/p$ if $q > p$, we have $D^{t_1} \in (H^p, H^q)$ (this is obvious if $p \leq q$, while the $p > q$ case is the Hardy-Littlewood theorem on fractional integrals; see, for example [14, Theorem 9]). If p, q are as hypothesised, it is readily verified that t_1 is larger than the index t_0 of Theorem 4.3, and so (H^p, H^q) is not solid. \square

For $0 < p < 2 < q < \infty$, it was proved by the third author [V, Theorem 2] that $\lambda = (\lambda_n) \in (A^p, A^q)$ if $\lambda_n = O(n^{2/q-2/p})$, and that this index is best possible. We now recover this result by applying Corollary 1.4 (or Theorem 1.3) to Lemma 4.1. More generally, we shall get separate necessary and sufficient conditions for a sequence to be in (A_α^p, A_β^q) for all allowable values of p, q, α, β . We begin with the sufficient conditions.

Theorem 4.5. *Suppose $p, q > 0$, and $\alpha, \beta > -1$. Let $p_1 = \max\{p, 2\}$, $q_1 = \min\{q, 2\}$, $s = q_1^{-1} - p_1^{-1}$, and $r = (2 + \beta)q^{-1} - (2 + \alpha)p^{-1} - s$. If*

$$(n^{-r} \lambda_n) \in l(\infty, s^{-1}) \quad (4.6)$$

then $\lambda = (\lambda_n) \in (A_\alpha^p, A_\beta^q)$. The index r is best possible in all cases.

Proof. The case $0 < q \leq 2 \leq p < \infty$ is one direction of Lemma 4.1. Let us look at the case $0 < p < 2 < q < \infty$. By Theorem 1.3, $I^{t_1} \in (A_\alpha^p, A_\alpha^2)$ and $I^{t_2} \in (A_\beta^2, A_\beta^q)$, where $t_1 = (2 + \alpha)(p^{-1} - 2^{-1})$ and $t_2 = (2 + \beta)(2^{-1} - q^{-1})$. By a (trivial) special case of Lemma 4.1, $\mu_n \in (A_\alpha^2, A_\beta^2)$ if and only if $(n^{(\alpha-\beta)/2} \mu_n) \in l^\infty$. Putting everything together, we get that $(\lambda_n) = (n^{-t_1-t_2} \mu_n) \in (A_\alpha^p, A_\beta^q)$ if $n^{t_1+t_2+(\alpha-\beta)/2} \lambda_n$ is bounded. Since

$$t_1 + t_2 + (\alpha - \beta)/2 = (2 + \alpha)p^{-1} - (2 + \beta)q^{-1},$$

and $s = 0$, we obtain (4.6).

In the case $0 < p, q < 2$, $\lambda \in (A_\alpha^p, A_\beta^q)$ if λ_n can be written as $n^{-t_1} \mu_n$, where t_1 is as before, and $(\mu_n) \in (A_\alpha^2, A_\beta^q)$; in the case $p, q > 2$, $\lambda \in (A_\alpha^p, A_\beta^q)$ if λ_n can be written as $n^{-t_2} \mu_n$, where t_2 is as before, and $(\mu_n) \in (A_\alpha^p, A_\beta^2)$. In both cases, (4.6) follows by applying Lemma 4.1 to (μ_n) .

It remains to show that r in (4.6) is maximal. By Corollary 1.4, we may assume without loss of generality that $\alpha = \beta = 0$. By an easy dyadic-block argument, it is equivalent to show that for every $\delta > 0$, there are sequences $(\lambda_n) = O(n^{r+\delta})$ not in (A^p, A^q) . For the case $0 < p < 2 < q < \infty$, it suffices to note that if $\delta > 0$ then $D^{-\delta+2/p-1} \in A^p$, but $D^r D^{-\delta+2/p-1} \notin A^q$ if $r \geq \delta + 2/q - 2/p$. Sharpness for $p < 2, q \leq 2$ and for $p > 2, q \geq 2$ follows from Theorem 4.2 (and conversely the sharpness of the index t_0 in Theorem 4.2 follows from this theorem). The case $p \geq 2 \geq q$ follows from Lemma 4.1. \square

We now state the corresponding necessary conditions for a sequence to be a multiplier (Theorems 4.7 and 4.8). A weaker version of 4.7 could be proved in a similar fashion to Theorem 4.5 (with fractional differentiation replacing fractional integration), but lacunary sequences provide a simpler route to the full-strength result. Note that the exponent r in Theorem 4.7 is sharp when $p \geq 1$, but not when $p < 1$. In the latter case, Theorem 4.8 gives the sharp order of growth.

Theorem 4.7. *Suppose $p, q > 0$, $\alpha, \beta > -1$, $r = (1 + \beta)q^{-1} - (1 + \alpha)p^{-1}$, and $s = \max\{0, q^{-1} - p^{-1}\}$. If $\lambda = (\lambda_n) \in (A_\alpha^p, A_\beta^q)$ then $(n^{-r}\lambda_n) \in l(\infty, s^{-1})$. The converse is true if $p \geq 1$ and λ is lacunary.*

Proof. By Corollary 1.4, we may as well assume $\alpha = \beta = 0$. Let (a_k) be a lacunary sequence whose terms are zero except possibly at positions $n_k \in B_k$, $k \geq 0$. Writing $b_k = 2^{-k/p}a_{n_k}$, the assumption $(a_n) \in A^p$ is equivalent to $(b_k) \in l^p$. Similarly applying $\lambda \in (A^p, A^q)$ to (a_n) , we get $(b_k\lambda_{n_k}2^{k(1/p-1/q)}) \in l^q$. Since $n_k \in B_k$ is arbitrary, it follows that $(b_k\mu_k) \in l^q$ for every $(b_k) \in l^p$, where $\mu_k = 2^{k(1/p-1/q)} \max\{|\lambda_n| : n \in B_k\}$. The required necessity now follows from the well-known result that $(l^p, l^q) = l^{1/s}$.

For the partial converse, suppose λ is lacunary and $(n^{-r}\lambda_n) \in l(\infty, s^{-1})$. Thus $\lambda_n = 0$ unless $n = n_k \in B_k$ and $(n_k^{-r}\lambda_{n_k}) \in l^{1/s}$. The Cauchy integral formula implies that if $f = (b_n) \in H^p$, then $|b_n| \leq \|f\|_{H^p}$ for all $p \geq 1$. This and (2.2) imply that if $(a_n) \in A^p$ then $(n_k^{-1/p}a_{n_k}) \in l^p$. Since $(l^p, l^q) = l^{1/s}$, the required result follows. \square

Theorem 4.8. *Suppose $0 < p < 1$, $0 < q < \infty$, and $\alpha, \beta > -1$. Let*

$$r = \begin{cases} \frac{1+\beta}{q} - \frac{2+\alpha}{p} + 1, & \text{if } 1 \leq q \\ \frac{2+\beta}{q} - \frac{2+\alpha}{p}, & \text{if } p < q < 1 \\ \frac{1+\beta}{q} - \frac{1+\alpha}{p}, & \text{if } q \leq p. \end{cases}$$

If $\lambda = (\lambda_n) \in (A_\alpha^p, A_\beta^q)$ then $\lambda_n = O(n^r)$. Furthermore, given any values of p, q, α, β , the number r is minimal in the last statement.

Proof. As usual, we may assume $\alpha = \beta = 0$. Theorem 4.7 implies (a sharper version of) the case $q \leq p$, so we need only look at the other two cases. By Lemma 3.8, $\lambda D^s \in A^q$ for each $s < 2/p - 1$, $\lambda \in (A^p, A^q)$, and so by parts (i) and (ii) of Lemma 1.2, we must have $\lambda_n = O(n^t)$ for any $t > r$. This is close to what we need but we need to argue a little more delicately to get $t = r$.

We first note that $(A^p, l^\infty) = \{(\lambda_n) : n^{2/p-1}\lambda_n \in l^\infty\}$, whenever $0 < p \leq 1$. This is a special case of a result of Blasco [6, 4.5] (alternatively note that containment in one direction follows from Lemma 1.2(i), and the opposite direction follows from the fact that $H^{p/2} \subset A^p$ and the corresponding result for H^p in [10, Theorem 6.5]). It follows that if $0 < 1/d_n \neq O(n^{1-2/p})$ then there exists $(a_n) \in A^p$ such that $a_n \neq O(d_n)$ (otherwise $(1/d_n) \in (A^p, l^\infty)$). Suppose therefore that $\lambda_n \neq O(n^r)$ and so there exist a sequence of integers n_k such that $\lambda_{n_k} > k^2 n_k^r$. Choosing $(a_n) \in A^p$ such that $k|a_{n_k}| > n_k^{2/p-1}$, we see that $(\lambda_n a_n)$ cannot be in A^q since it violates 1.2(i) ($q < 1$) or 1.2(ii) ($1 \leq q$).

We are left with proving minimality of r . In the case $q \geq 1$, simply consider lacunary multipliers and use Lemma 1.2(i). Parts (a) and (b) of Corollary 1.4 imply r is minimal when $0 < p \leq q < 1$ and when $0 < q < p < 1$, respectively. \square

We now compare our L^∞ -type necessary/sufficient multiplier estimates with the L^q -type estimates for multipliers between unweighted Bergman spaces of the third author [29]. Our sufficient conditions are easily seen to imply the following result.

Corollary 4.9. *In any of the following circumstances, $\lambda = (\lambda_n) \in (A^p, A^q)$.*

- (i) $0 < p, q \leq 2$ and $\sum_{n=1}^{\infty} n^{-1-q/2+2q/p} |\lambda_n|^q < \infty$.
- (ii) $p, q \geq 2$ and $\sum_{n=1}^{\infty} n^{-2+q/2+q/p} |\lambda_n|^q < \infty$.
- (iii) $p \leq 2 \leq q$ and $\sum_{n=1}^{\infty} n^{-2+2q/p} |\lambda_n|^q < \infty$.
- (iv) $q \leq 2 \leq p$ and $\sum_{n=1}^{\infty} n^{-1+q/p} |\lambda_n|^q < \infty$.

Proposition 1 of [29] also gives a sufficient condition for each of four cases, but the powers of n in the sums are different and they change at $p = 1$ rather than $p = 2$ (see [29] for details). Naturally, one can also use the L^q estimates of [29] to derive L^∞ estimates. Doing so, one sees that the two sets of results are incomparable if $0 < p \leq 1, q > 0$, and also if $1 < p < 2, q < p'$. However, Theorem 4.5 implies Proposition 1 when $1 < p \leq 2 \leq p' < q$ and when $p > 2, q > 0$; in fact if $q > \max(2, p')$, then the condition in Corollary 4.9 has a strictly smaller power of n than its counterpart in Proposition 1.

We next examine L^q necessary conditions to be a multiplier. Since it is routine to derive such conditions from Theorems 4.7/4.8, we state a result only for the case where we get better estimates than those of the corresponding Theorem 8 of [29]; we use “ $O(N)$ ” format for greater ease of comparability with Theorem 8.

Corollary 4.10. *Suppose $\lambda = (\lambda_n) \in (A^p, A^q)$, and that either $0 < q \leq p$ or $p \geq 1, q > 0$. Then $\sum_{n=1}^N n^{q/p-1-\epsilon} |\lambda_n|^q = O(N)$ for every $\epsilon > 0$.*

Corollary 4.10 is stronger than Theorem 8 when $q < \min(2, p)$ and also when $1 < p < 2, q > p'$. In all other cases where it is applicable it is weaker; in fact, Theorems 8 and 4.7/4.8 are incomparable. In cases not handled by Corollary 4.10, Theorems 8 and 4.8 are also incomparable except when $0 < p < 1, q \geq 2$. Theorem 8 states that if $\lambda \in (A^p, A^q)$ for some $0 < p < \infty, 2 \leq q < \infty$, then $\sum_{n=1}^N n^{2q/p-q} |\lambda_n|^q = O(N)$. This implies that (A^p, A^q) multipliers ($q \geq 2$) have terms which are $o(n^{1/q+1-2/p})$, a sharper version of Theorem 4.8 for $0 < p < 1, q \geq 2$.

§5. Univalent Bergman Functions

The integrability of derivatives of conformal maps is one of the central questions in function theory. We show here (at least in the unweighted case) that the gap in Corollary 1.4 between fractional integration and differentiation (parts (a) and (b)) can be bridged “to within ϵ ” if the function is univalent. Below, \mathcal{U} is the class of functions univalent on \mathbf{D} , and $S \subset \mathcal{U}$ denotes the normalised univalent functions that satisfy $f(0) = 0$ and $f'(0) = 1$.

Proposition 5.1. *If $f \in A^p \cap \mathcal{U}$ and $0 < t \leq 1$ then $D^t f \in A^q$ for all $q^{-1} > p^{-1} + t/2$.*

Proof. It was shown in [20] that if a quasiconformal mapping on the unit ball $B \subset \mathbf{R}^n$ is in $L^p(B)$, then its derivative is in $L^q(B)$ for all $q^{-1} > p^{-1} + 1/n$. Since univalent mappings are quasiconformal, the $n = 2$ case gives the $t = 1$ case of our result. The case $0 < t < 1$ then follows easily from the $t = 1$ case, the identity $D^t = D^1 * I^{1-t}$, and Theorem 1.3(a). \square

A theorem of Prawitz [10, Theorem 3.16] says that $\mathcal{U} \subset H^p$ for all $p < 1/2$, and hence $\mathcal{U} \subset A^p$ for all $p < 1$ (the Koebe function shows that a univalent function need not be in A^1). Thus the above proposition implies that $D^t f \in A^q$ for all $q < 2/(2 + t)$. Alternatively, this fact follows in the case $t = 1$ from (the less subtle part of) a well-known result of Brennan [7] which says that if $f \in S$ then $|f'|^q$ is integrable on the unit disc for all $-1 - \epsilon < q < 2/3$ as long as $\epsilon > 0$ is suitably small; the case $0 < t < 1$ then follows as before.

Proposition 5.2. *If $f \in \mathcal{U}$ and $0 < t \leq 1$ then $D^t f \in A^q$, for any $q < 2/(t + 2)$. The statement is false for $q = 2/(t + 2)$.*

Proof. The first statement has been proved in the above paragraph, so we need only show that the exponent $q = 2/(t + 2)$ cannot be achieved. For $t = 1$ (i.e., $q = 2/3$), an explicit calculation shows that the Koebe function $k(z) = z/(1 - z)^2$ is a counterexample. For $0 < t < 1$, we have $D^t k = D^{1+t}$, and so $D^t k \notin A^{2/(t+2)}$ by Lemma 3.8. \square

References

- [1] P. AHERN and M. JEVIĆ. Duality and multipliers for mixed norm spaces. *Michigan Math. J.* **30** (1983), 53–64.
- [2] J.M. ANDERSON. Coefficient multipliers and solid spaces. *J. Analysis* **1** (1993), 13–19.
- [3] J.M. ANDERSON and D.GIRELA. Inequalities of Littlewood-Paley type, multipliers and radial growth of the derivative of analytic functions. *J. reine angew. Math.* **465** (1995), 11–40.
- [4] J.M. ANDERSON and A.L. SHIELDS. Coefficient multipliers of Bloch functions. *Trans. Amer. Math. Soc.* **224** (1976), 255–265.
- [5] S. AXLER. Bergman spaces and their operators. In *Survey of Some Recent Results in Operator Theory*, Vol. 1 (edited by J.B. CONWAY and B.B. MORREL), Pitman Research Notes in Mathematics, vol. **171** (Longman Scientific & Technical, Harlow, Essex, 1988), pp. 1–50.
- [6] O. BLASCO. Multipliers on spaces of analytic functions. *Can. J. Math.* **47** (1995), 44–64.
- [7] J.E. BRENNAN. The integrability of the derivative in conformal mapping. *J. London Math. Soc.* **18** (1978), 261–272.
- [8] S. BUCKLEY, M.S. RAMANUJAN, and D. VUKOTIĆ. Bounded and compact coefficient multipliers between Bergman and Hardy spaces. In preparation.

- [9] D. CAMPBELL and R. LEACH. A survey of H^p multipliers as related to classical function theory. *Complex Variables* **3** (1983), 85–111.
- [10] P.L. DUREN. *Theory of H^p spaces*. (Academic Press, New York, 1970).
- [11] P.L. DUREN and A.L. SHIELDS. Coefficient multipliers of H^p and B^p spaces. *Pacific J. Math.* **32** (1970), 69–78.
- [12] P.L. DUREN and G.D. TAYLOR. Mean growth and coefficients of H^p functions. *Illinois J. Math.* **14** (1970), 419–423.
- [13] T. FLETT. Some theorems on fractional integrals. *Proc. Camb. Phil. Soc.* **55** (1959), 31–50.
- [14] T. FLETT. Mean values of power series. *Pacific J. Math.* **25** (1968), 463–494.
- [15] T. FLETT. The dual of an inequality of Hardy and Littlewood and some related inequalities. *J. Math. Anal. Appl.* **38** (1972), 746–765.
- [16] G.H. HARDY and J.E. LITTLEWOOD. Some properties of fractional integrals, II. *Math. Z.* **34** (1932), 403–439.
- [17] C.A. HOROWITZ. Zeros of functions in the Bergman spaces. *Duke. J. Math.* **41** (1974), 693–710.
- [18] M. JEVTIĆ and I. JOVANOVIĆ. Coefficient multipliers of mixed norm spaces, *Canad. Math. Bull.* **36** (1993), 283–285.
- [19] M. JEVTIĆ and M. PAVLOVIĆ. On multipliers from H^p to l^q , $0 < q < p < 1$. *Arch. Math.* **56** (1991), 174–180.
- [20] P. KOSKELA. An inverse Sobolev lemma. *Rev. Mat. Iberoamericana* **10** (1994), 123–141.
- [21] C.N. KELLOGG. An extension of the Hausdorff-Young theorem. *Michigan Math. J.* **18** (1971), 121–127.
- [22] D. LUECKING. Embedding theorems for spaces of analytic functions via Khinchine’s inequality. *Michigan Math. J.* **40** (1993), 333–358.
- [23] J.E. LITTLEWOOD and R.E.A.C. PALEY. Theorems on Fourier series and power series (II). *J. London Math. Soc.* **42** (1931), 52–89.
- [24] J. MARSDEN. *Basic Complex Analysis*. (W.H. Freeman, San Francisco, 1972).
- [25] M. MATELJEVIĆ and P. PAVLOVIĆ. L^p behaviour of the integral means of analytic functions. *Studia Math.* **77** (1984), 219–237.
- [26] M. MATELJEVIĆ and M. PAVLOVIĆ. Multipliers of H^p and BMOA. *Pacific J. Math.* **146** (1990), 71–84.
- [27] A. NAKAMURA, F. OHYA, and H. WATANABE. On some properties of functions in weighted Bergman spaces. *Proc. Fac. Sci. Tokai Univ.* **15** (1979), 33–44.
- [28] M. PAVLOVIĆ. Mixed norm spaces of analytic and harmonic functions II. *Publ. Inst. Math. Beograd* **41** (55) (1987), 97–110.
- [29] D. VUKOTIĆ, On the coefficient multipliers of Bergman spaces. *J. London Math. Soc.* **50** (1994), 341–348.
- [30] P. WOJTASZCZYK. On multipliers into Bergman spaces and Nevanlinna class. *Canadian Math. Bull.* **33** (1990), 151–161.

- [31] A. ZABULIONIS. Differential operator in spaces of analytic functions. *Liet. Mat. Rinkinys* **24** (1984), 53–59. (English translation in: *Lithuanian Math. J.* 1984, 32–36).
- [32] K. ZHU. Duality of Bloch spaces and norm convergence of Taylor series. *Michigan Math. J.* **38** (1991), 89–101.

S. Buckley: Department of Mathematics, Maynooth College, Maynooth, Co. Kildare, Ireland (e-mail: sbuckley@maths.may.ie).

P. Koskela: Department of Mathematics, University of Jyväskylä, P.O.Box 35, Fin-40351 Jyväskylä, Finland (e-mail: pkoskela@math.jyu.fi).

D. Vukotić: Departamento de Matemáticas, Universidad Autónoma de Madrid, Ciudad Universitaria de Cantoblanco, 28049 Madrid, Spain (e-mail: dragan.vukotic@uam.es, dragan@bosque.sdi.uam.es.)