

LEAVING CERTIFICATE EXAMINATION, 1976

APPLIED MATHEMATICS - HIGHER LEVEL

FRIDAY, 25 JUNE - MORNING, 9.30 to 12


Six questions to be answered. All questions carry equal marks.

Mathematics Tables may be obtained from the Superintendent.

Take the value of g to be 9.8 metres/second². \vec{i} and \vec{j} are perpendicular unit vectors.

1. Show that, if a particle is moving in a straight line with constant acceleration k and initial speed u , the distance travelled in time t is given by $s = ut + \frac{1}{2}kt^2$. Two points a and b are a distance l apart. A particle starts from a and moves towards b in a straight line with initial velocity u and constant acceleration k . A second particle starts at the same time from b and moves towards a with initial velocity $2u$ and constant deceleration k . Find the time in terms of u, l at which the particles collide, and the condition satisfied by u, k, l if this occurs before the second particle returns to b .
2. A particle is projected upwards with a speed of 35 m/s from a point O on a plane inclined at 45° to the horizontal. The plane of projection meets the inclined plane in a line of greatest slope and the angle of projection, measured to the inclined plane, is ϕ . Write down the velocity of the particle and its displacement from O , in terms of \vec{i} and \vec{j} , after time t seconds. If the particle is moving horizontally when it strikes the plane at q prove that $\cot \phi = 3$ and calculate $|oq|$.

3. The diagram shows a light inelastic string, passing over a fixed pulley B , connecting a particle A of mass $3M$ to a light movable pulley C . Over this pulley passes a second light inelastic string to the ends of which are attached particles D, E of masses $2M, M$ respectively. Show in separate diagrams the forces acting on A, D and E . Write down the three equations of motion involving the tensions T, S in the strings, the acceleration of A and the common acceleration of D, E relative to C . Show that $T = 2S = 48Mg/17$.



4. A light smooth ring of mass M is threaded on a smooth fixed vertical wire and is connected by a light inelastic string, passing over a fixed smooth peg at a distance l from the wire, to a particle of mass $2M$ hanging freely. The system is released from rest when the string is horizontal. Explain why the conservation of energy can be applied to the system. If the ring descends a distance of x while the particle rises through a distance y show that

$$x^2 = y^2 + 2ly \text{ and } (l + y)\dot{y} = x\dot{x}$$

where $\dot{x} = \frac{dx}{dt}$, $\dot{y} = \frac{dy}{dt}$ are the speeds of the ring and particle respectively. Find \dot{x} when (i) $x = l$ and

(ii) when $x = \frac{4l}{3}$.

5. State the laws governing the oblique collision of elastic spheres.

A sphere of mass M moving with speed u collides obliquely with a second smooth sphere at rest. The direction of motion of the moving sphere is inclined at 45° to the line of centres at impact, and the coefficient of restitution is $\frac{1}{2}$. After impact the directions of motion of the spheres are at right angles.

Find the mass of the second sphere in terms of M , and the velocities of the two spheres after impact in terms of u . Hence show that one quarter of the kinetic energy is lost.

6. Two uniform rods ab , bc of lengths $2l$, $2r$ and of weights $2W$, $3W$ respectively are smoothly hinged together at b . They stand in equilibrium in a vertical plane with the end a resting on rough horizontal ground and the end c resting against a smooth vertical wall. The point a is farther from the wall than b and the rods ab , bc are inclined at angles α , 45° respectively to the horizontal where $\alpha > 45^\circ$. Show in separate diagrams the forces acting on each rod. By considering separately the equilibrium of the system abc and of the rod bc , find the coefficient of friction at a and show that $\tan \alpha = \frac{8}{3}$.

7. Define simple harmonic motion.

A particle of mass 2 kg is attached to the ends of two light elastic strings, each of natural length 1 m and elastic constant 49 N/m. The other ends of the two strings are attached to two fixed points a and b in the same vertical line, where a is 4 m above b . The particle is released from rest from the midpoint of ab . By considering the forces acting on the particle when it is x metres from a , where $2 < x < 2.4$, show that it is moving with simple harmonic motion. Find the least time taken for the particle to reach the point $x = 2.3$, and find its speed there.

8. A pendulum of a clock consists of a thin uniform rod ab of mass M and length $6l$ to which is rigidly attached a uniform circular disc of mass $4M$ and radius l with the centre of the disc being at the point c on ab where $bc = l$. Using the parallel axes theorem for the disc, show that the moment of inertia of the pendulum about an axis at a perpendicular to the plane of the disc is $114Ml^2$.

The pendulum is free to oscillate in a vertical plane about such a fixed horizontal axis at a . It is released from rest with ab horizontal. Find the speed of b when ab is vertical.

9. An atomic nucleus of mass M is repelled from a fixed point o by a force Mk^2x^{-5} , where x is the distance of the nucleus from o and k is a constant. It is projected directly towards o with speed $\frac{2k\sqrt{3}}{d^2}$ from a point a where $|oa| = d$. Find the speed of the nucleus when it reaches the midpoint of oa and find how near it gets to o .

10. (i) Using Taylor's theorem (Mathematics Tables, p.42) find the first two terms in the Taylor series for e^{x^2} in the neighbourhood of $x = 0$, i.e. the Maclaurin series for e^{x^2} .

(ii) State Archimedes principle for a body wholly or partly immersed in a liquid.

A uniform thin rod is of length $2a$, of weight $4W$ and specific gravity $\frac{4}{9}$. The rod rests in equilibrium in an inclined position partly immersed in water with its lower end freely pivoted to a fixed point at a depth $\frac{2a}{3}$ below the surface of the water. Show in a diagram the forces acting on the rod and calculate the inclination of the rod to the vertical.