


APPLIED MATHEMATICS – HIGHER LEVEL

FRIDAY, 17 JUNE – MORNING, 9.30 – 12.00

Six questions to be answered. All questions carry equal marks.
 Mathematics Tables may be obtained from the Superintendent.
 Take the value of g to be 9.8 m/s^2 .
 Marks may be lost if all your work is not shown or you do not indicate where a calculator has been used.


1. (a) A particle moving in a straight line with uniform acceleration describes 23 m in the fifth second of its motion and 31 m in the seventh second. Calculate its initial velocity.
- (b) A particle falls freely from rest from a point o , passing three points a , b and c , the distances ab and bc being equal. If the particle takes 3 s to pass from a to b and 2 s from b to c , calculate $|ab|$.
2. (a) Two boats move with constant speed 5 m/s relative to the water and both cross a straight river of width 72 m flowing with constant speed 3 m/s parallel to the banks. One crosses by the shortest path and the other in the shortest time. Show that the difference in the times taken is 3.6 s.
- (b) Two ships A and B move with constant speeds $2u$ and u respectively. At a certain instant, B is 2400 m due east of A and moving northwards. Show that A must move in the direction 30° North of East in order to intercept B and find (in terms of u) the time it takes to intercept B .
3. (a) A particle which is projected with speed u has a horizontal range $\frac{3u^2}{49}$. Calculate the two possible angles of projection.
- (b) A particle is projected up an inclined plane with initial speed $13u$. The line of projection makes an angle $\tan^{-1}(\frac{5}{12})$ with the plane and the plane is inclined at 45° to the horizontal. (The plane of projection is vertical and contains the line of greatest slope.) The particle strikes the plane at a point p . If the coefficient of restitution between the particle and the plane is 0.4, show that the particle rises vertically from p and strikes p again on the second bounce.

4. One end of a light inextensible string is attached to a mass of 6 kg which rests on a rough horizontal table. The coefficient of friction between the mass and the table is $\frac{1}{6}$. The string passes over a smooth fixed pulley at the edge. Then it passes under a smooth movable pulley of mass 4 kg and over a smooth fixed pulley. A mass of 2 kg is attached to its other end.



- (i) Show on separate diagrams the forces acting on each mass.
- (ii) Calculate the acceleration of each mass and the tension in the string in terms of g , the acceleration due to gravity.


5. Two smooth spheres, A and B , of equal radii, have masses 4 kg and 8 kg respectively. They lie at rest on a smooth horizontal floor so that the line joining their centres is perpendicular to the vertical wall.



A is projected towards B with speed u and collides with B . B then hits the wall, rebounds and collides with A again. This final collision reduces B to rest. If the coefficient of restitution between A and B is $\frac{1}{4}$, calculate

- the coefficient of restitution between B and the wall.
 - the final velocity of A in terms of u .
 - the total loss of energy due to the three collisions.
6. A particle of mass 8 kg is describing a circle, with constant speed v , on a smooth horizontal table. It is connected by a light inextensible string of length 3 m to a point which is 1 m vertically above the centre of the circle.
- Calculate the tension in the string.
 - Show that the particle will remain in contact with the table if $v \leq \sqrt{8g}$.
 - If the speed of the particle is increased to $\sqrt{9.1g}$, calculate the height at which the particle rotates above the table.

7. Two equal uniform rods ab and bc each of length $2l$ and weight W , are freely jointed at b and rest in equilibrium, in a vertical plane, across two smooth horizontal pegs at the same horizontal level and distant $\frac{16l}{27}$ apart.




- Show in separate diagrams the forces acting on each rod.
 - Show that the inclination of each rod to the vertical is $\sin^{-1}(\frac{2}{3})$.
 - Determine the magnitude and direction of the reaction at b .
8. Show that the moment of inertia of a uniform rod of mass m and length $2l$, about an axis through its centre of mass perpendicular to the rod is $\frac{1}{3}ml^2$.
Three of these rods are joined together at their ends to form a triangle abc . The triangle is free to rotate about a fixed horizontal axis through a , perpendicular to its plane. Find the period of small oscillations about the equilibrium position.

9. State the Principle of Archimedes.

A uniform rod of weight W and of length $2l$, in equilibrium, is supported at one end by a vertical force T and is immersed in water as shown in the diagram.

The relative density of the rod is $\frac{7}{16}$.



- Calculate the length of the immersed part of the rod.
- Show that $T = \frac{3W}{7}$.

10. (a) Solve the differential equation

$$\frac{dx}{dt} = \sqrt{100 - 4x^2}$$

if $x = 5$ when $t = 0$.

- (b) A particle of mass m is projected vertically upwards with speed 120 m/s in a medium where there is a resistance of $0.098v^2$ per unit mass of the particle when v is the speed. Calculate the time taken to reach the highest point.