

CARTER-PAYNE HOMOMORPHISMS AND BRANCHING RULES FOR ENDOMORPHISM RINGS OF SPECHT MODULES

HARALD ELLERS AND JOHN MURRAY

ABSTRACT. Let Σ_n be the symmetric group of degree n , and let F be a field of characteristic $p \neq 2$. Suppose that λ is a partition of $n+1$, that α and β are partitions of n that can be obtained by removing a node of the same residue from λ , and that α dominates β . Let S^α and S^β be the Specht modules, defined over F , corresponding to α , respectively β . We give a very simple description of a non-zero homomorphism $\theta : S^\alpha \rightarrow S^\beta$ and present a combinatorial proof of the fact that $\dim \operatorname{Hom}_{F\Sigma_n}(S^\alpha, S^\beta) = 1$. As an application, we describe completely the structure of the ring $\operatorname{End}_{F\Sigma_n}(S^\lambda \downarrow_{\Sigma_n})$. Our methods furnish a lower bound for the Jantzen submodule of S^β that contains the image of θ .

1. INTRODUCTION

Let n be a positive integer and let Σ_n denote the symmetric group of degree n . For any ring R and any partition α of n , the Specht module S_R^α is defined to be the submodule of the permutation module $R\Sigma_n \uparrow^{\Sigma_n}$ spanned by certain elements called polytabloids, where Σ_α is the Young subgroup associated to α and $R\Sigma_\alpha$ is the trivial $R\Sigma_\alpha$ -module. (See [4] for definitions.) Specht modules play a central role in the representation theory of the symmetric group. Now suppose that F is a field. If F has characteristic 0, the Specht modules defined over F are the simple $F\Sigma_n$ -modules; if F has characteristic p the heads of the Specht modules S_F^α , with α p -regular, are the simple $F\Sigma_n$ -modules.

It would be useful to understand $\operatorname{Hom}_{F\Sigma_n}(S^\alpha, S^\beta)$ for all partitions α and β of n when F has positive characteristic, but this problem has not been solved in general. In [2], Carter and Payne exhibited some types of partitions α and β for which $\operatorname{Hom}_{\Sigma_n}(S^\alpha, S^\beta) \neq 0$ even though $\alpha \neq \beta$, generalizing earlier results of Carter and Lusztig [1].

Among the examples discovered by Carter and Payne are *one-box shift* homomorphisms. Suppose that the Young diagram $[\beta]$ of β can be obtained by moving a single node in the diagram $[\alpha]$; then we say that β is *one-box-shift* of α (see (5)). This is equivalent to saying that there exists a partition λ of $n+1$ such that α and β can each be obtained by removing a node from the Young diagram of $[\lambda]$. Carter and Payne showed that if the node in $[\alpha] \setminus [\beta]$ has the same p -residue as the node in $[\beta] \setminus [\alpha]$, and α dominates β , then $\operatorname{Hom}_{\Sigma_n}(S^\alpha, S^\beta) \neq 0$ (the *residue* of the node in the i -th row and j -th column of a Young diagram is the integer $j - i$; the p -residue is just $j - i$ taken modulo p).

In fact, in this case $\dim \operatorname{Hom}_{F\Sigma_n}(S^\alpha, S^\beta) = 1$, when the characteristic of F is greater than 2. This was proved in the context of Iwahori-Hecke algebras by Lyle

Date: January 30, 2009.

1991 Mathematics Subject Classification. 20C20, 20C30.

Key words and phrases. one-box-shift homomorphism, Jucys-Murphy elements.

[8]. For the group algebras, this result seems to have been known to experts in the area for some time. For example, A. Kleshchev claims, apparently incorrectly, in [5, p498], that this is a special case of the main result in [1]. However, we do not believe that a complete proof of this theorem had been published before Lyle's paper.

Let L_{n+1} be the Jucys-Murphy element consisting of the sum in $F\Sigma_{n+1}$ of all transpositions in Σ_{n+1} that are not in Σ_n . We would like to point out that results from [3] make it possible to give the following simple description of one-box shift homomorphisms:

Theorem 1.1. *Let F be a field of characteristic $p > 0$ and let λ be a partition of $n + 1$. Suppose that α and β can each be obtained by removing a node from λ , with $\alpha \triangleright \beta$, and that the removed nodes have the same p -residues. Identify S_F^α and S_F^β as subquotients of $S_F^\lambda \downarrow_{\Sigma_n}$. Then multiplication by an appropriate polynomial in L_{n+1} induces a non-zero homomorphism from S_F^α to S_F^β .*

Proof. Assume that λ has t removable nodes, with p -residues r_1, \dots, r_t . Let λ_i be the partition of n obtained by removing the i -th removable node from $[\lambda]$. Following James [4, 9.3], let S_i^λ be the subspace of S_F^λ spanned by those polytabloids e_t where t is a standard λ -tableau such that the symbol $n + 1$ is in one of the first i removable nodes of t . Then S_i^λ is a Σ_n -submodule of S^λ , and $S_i^\lambda/S_{i-1}^\lambda \cong S^{\lambda_i}$. The Σ_n -series

$$0 \subset S_0^\lambda \subset S_2^\lambda \subset \dots \subset S_t^\lambda = S^\lambda$$

is called the Specht series of $S_F^\lambda \downarrow_{\Sigma_n}$.

Suppose that $\alpha = \lambda_u$ and that $\beta = \lambda_v$. Then $u < v$. In [3], we showed that $S_u^\lambda \prod_{i=u}^{v-1} (L_{n+1} - r_i) \subseteq S_v^\lambda$ and that this map induces a well-defined nonzero homomorphism $S_F^\alpha = S_u^\lambda/S_{u-1}^\lambda \rightarrow S_v^\lambda/S_{v-1}^\lambda = S_F^\beta$.

If $p > 2$, then $\text{Hom}_{\Sigma_n}(S_F^\alpha, S_F^\beta)$ is 1-dimensional, and this map must be a nonzero multiple of the homomorphism defined in [2]. \square

In order to effectively use this result, we need some information about the action of L_{n+1} on S^λ . The following is a variant of (3.3) in [9]:

Lemma 1.2. *Let λ be a partition of $n + 1$ and let t be a λ -tableau (not necessarily standard). Suppose that $n + 1$ occupies a column of length r in t . Then λ has one removable node (r, c) for some $c \geq 1$. Let R be the set of symbols in t whose columns have length strictly smaller than r . Then in $S_{\mathbb{Z}}^\lambda$ we have*

$$e_t L_{n+1} = (c - r)e_t + \sum_{i \in R} e_t(n + 1, i).$$

For example, let $\lambda = [4, 3, 1]$, $n = 7$ and $t = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & 7 & \\ \hline 8 & & & \\ \hline \end{array}$. Then $n + 1 = 8$ occupies a node of residue -2 in t . We use Lemma 1.2 to compute $e_t(L_8 + 2)$. For emphasis we replace the symbol $n + 1$ by \bullet . The above lemma show that:

$$(1) \quad e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & 7 & \\ \hline \bullet & & & \\ \hline \end{array}}(L_8 + 2) = e_{\begin{array}{|c|c|c|c|} \hline 1 & \bullet & 3 & 4 \\ \hline 5 & 6 & 7 & \\ \hline 2 & & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & \bullet & 7 & \\ \hline 6 & & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & \bullet & 4 \\ \hline 5 & 6 & 7 & \\ \hline 3 & & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & \bullet & \\ \hline 7 & & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & \bullet \\ \hline 5 & 6 & 7 & \\ \hline 4 & & & \\ \hline \end{array}}$$

We use this to illustrate Theorem 1.1. Let $\alpha = [4, 3]$ and $\beta = [3, 3, 1]$. For $p = 5$ and $n = 7$ there is a one-box shift $\theta : S^\alpha \rightarrow S^\beta$. Now $u = 1, v = 3$ and $r_1 = 3, r_2 = 1$ and $r_3 = -2$. The relevant polynomial in L_{n+1} is $(L_8 + 2)(L_8 - 1)$.

We find the image of the polytabloid e_t with $t = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & 7 & \\ \hline \bullet & & & \\ \hline \end{array}$ under θ . We again replace 8 by \bullet . First note that $e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & 7 & \\ \hline \bullet & & & \\ \hline \end{array}}$ is an element of S_3^λ whose image is $e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & 7 & \\ \hline \bullet & & & \\ \hline \end{array}}$, under $S_3^\lambda/S_2^\lambda \cong S^\alpha$. We consider the summands in $e_t(L_8 + 2)$ given in (1) in turn:

$$\begin{aligned} e_{\begin{array}{|c|c|c|c|} \hline 1 & \bullet & 3 & 4 \\ \hline 5 & 6 & 7 & \\ \hline 2 & & & \\ \hline \end{array}} (L_8 - 1) &= e_{\begin{array}{|c|c|c|c|} \hline 1 & 4 & 3 & \bullet \\ \hline 5 & 6 & 7 & \\ \hline 2 & & & \\ \hline \end{array}} \\ e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & \bullet & 7 & \\ \hline 6 & & & \\ \hline \end{array}} (L_8 - 1) &= e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & \bullet \\ \hline 5 & 4 & 7 & \\ \hline 6 & & & \\ \hline \end{array}} \\ e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & \bullet & 4 \\ \hline 5 & 6 & 7 & \\ \hline 3 & & & \\ \hline \end{array}} (L_8 - 1) &= e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 4 & \bullet \\ \hline 5 & 6 & 7 & \\ \hline 3 & & & \\ \hline \end{array}} \\ e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & \bullet & \\ \hline 7 & & & \\ \hline \end{array}} (L_8 - 1) &= e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & \bullet \\ \hline 5 & 6 & 4 & \\ \hline 7 & & & \\ \hline \end{array}} \\ e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & \bullet \\ \hline 5 & 6 & 7 & \\ \hline 4 & & & \\ \hline \end{array}} (L_8 - 1) &= e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & \bullet \\ \hline 5 & 6 & 7 & \\ \hline 4 & & & \\ \hline \end{array}} (3 - 1). \end{aligned}$$

Deleting \bullet from each tableau, we get a tableau of shape β . So

$$(2) \quad e_{\begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & 7 & \\ \hline \end{array}} \theta = e_{\begin{array}{|c|c|c|} \hline 1 & 4 & 3 \\ \hline 5 & 6 & 7 \\ \hline 2 & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 5 & 4 & 7 \\ \hline 6 & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 5 & 6 & 7 \\ \hline 3 & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 5 & 6 & 4 \\ \hline 7 & & \\ \hline \end{array}} + 2e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 5 & 6 & 7 \\ \hline 4 & & \\ \hline \end{array}}.$$

Our second goal is to complete our analysis (see [3]) of the endomorphism ring of the restriction of the Specht module S^λ to Σ_n .

Theorem 1.3. *Let F be a field of characteristic p not equal to 2 and let λ be a partition of $n + 1$. Then the blocks of $\text{End}_{F\Sigma_n}(S^\lambda \downarrow_{\Sigma_n})$ are parametrized by the p -residues of the removable nodes in $[\lambda]$. Let r denote the p -residue of a removable node of λ . Then the corresponding block of $\text{End}_{F\Sigma_n}(S^\lambda \downarrow_{\Sigma_n})$ has the structure of a truncated polynomial ring over F . Its dimension is equal to the number of removable nodes in $[\lambda]$ that have p -residue r .*

The corresponding statement for the induced Specht module is also true:

Theorem 1.4. *Let F be a field of characteristic p not equal to 2 and let λ be a partition of $n - 1$. Then the blocks of $\text{End}_{F\Sigma_n}(S^\lambda \uparrow_{\Sigma_n})$ are parametrized by the p -residues of the addable nodes in $[\lambda]$. Let r denote the p -residue of an addable node of λ . Then the corresponding block of $\text{End}_{F\Sigma_n}(S^\lambda \uparrow_{\Sigma_n})$ has the structure of a truncated polynomial ring over F . Its dimension is equal to the number of addable nodes in $[\lambda]$ that have p -residue r .*

It is interesting to compare these theorems to a result of Kleshchev, which says that if D is a simple $F\Sigma_n$ -module, then $\text{End}_{F\Sigma_{n-1}}(D \downarrow_{\Sigma_{n-1}})$ is also a direct sum of truncated polynomial algebras, with one summand for each block of $F\Sigma_{n-1}$. (See Theorems 11.2.7 and 11.2.8 of [6].)

In later sections, we revisit the dimension of the space of one-box-shift homomorphisms.

Theorem 1.5. *Let F be a field of characteristic not equal to 2 and let β be a one-box-shift of α . Then $\dim \text{Hom}_{F\Sigma_n}(S^\alpha, S^\beta) \leq 1$.*

Our aim is to show that a proof is possible that uses only the relatively elementary combinatorial tools developed by G. James in his monograph [4]. In particular, we do not use the Schur functor, nor do we make use of any of the deeper machinery

of algebraic groups. We also appear to avoid some technical difficulties that arise in the context of Iwahori-Hecke algebras c.f. [8].

All three theorems 1.3, 1.4 and 1.5 are false when $p = 2$. The proof of Theorem 1.5 relies on Lemma 2.1, which fails for $p = 2$. In the case of Theorems 1.3, and 1.4, when $p = 2$, we do not even know how to parametrize the blocks of $\text{End}_{F\Sigma_n}(S^\lambda \downarrow_{\Sigma_n})$ nor those of $\text{End}_{F\Sigma_n}(S^\lambda \uparrow^{\Sigma_n})$. For these reasons we now fix F as a field of characteristic $p \neq 2$.

The methods we use to prove Theorem 1.5 make it possible to explore how one-box-shift homomorphisms interact with Jantzen layers (see Section 7 for definitions).

Theorem 1.6. *Let λ be a partition of $n + 1$ and let F be a field of characteristic $p > 0$. Suppose that α and β can be obtained by removing a node of residue r_α and r_β , respectively, from λ , with $\alpha \succeq \beta$. Let $\theta : S_F^\alpha \rightarrow S_F^\beta$ be a Σ_n -homomorphism. If p^i divides $r_\alpha - r_\beta$ then $\theta(S_F^\alpha)$ is contained in the i -th Jantzen submodule of S_F^β .*

2. NOTATION AND BASIC RESULTS

For the rest of the paper, unless stated otherwise, F is a field of characteristic $p > 2$ and α is a fixed partition of n . We also fix a bijection

$$t : [\alpha] \rightarrow \{1, \dots, n\}.$$

The Young diagram $[\alpha]$ is a collection of boxes in the plane that is oriented left to right and top to bottom. This means that the first row is the one at the top and the first column is the one at the left. The (i, j) -th node of $[\alpha]$ is the box in the i -th row and the j -th column.

We refer to any map $T : [\alpha] \rightarrow \{1, \dots, n\}$ as an α -*tableau*. Informally, T is a filling of the nodes of $[\alpha]$, using elements from $\{1, \dots, n\}$, with repeats allowed. The value of T at a node $b = (r, c)$ is denoted by T_b or $T(r, c)$. If $i = T_b$, we say that node b of T *contains* the symbol i .

Identify Σ_n with the group of permutations of $\{1, \dots, n\}$. Permutations and homomorphisms will generally act on the right. For instance Σ_n acts on the set of all bijective α -tableau by functional composition. We can use the bijection t to give a contragredient action of Σ_n on all α -tableau: if $\pi \in \Sigma_n$ then define the α -tableau $T\pi$ as the map $T\pi = t \circ \pi^{-1} \circ t^{-1} \circ T$.

Suppose that α has r nonzero parts $[\alpha_1 \geq \alpha_2 \geq \dots \geq \alpha_r]$. An α -*tabloid* is an ordered partition of \hat{n} whose parts have cardinalities $\alpha_1, \alpha_2, \dots, \alpha_r$. For instance the sets $\{t_{ij} \mid j = 1, \dots, \alpha_i\}$ for $i = 1, \dots, r$, determine an α -tabloid that we shall denote by $\{t\}$.

Let R_t be the row stabilizer and let C_t be the column stabilizer of t . Denote by M^α the $F\Sigma_n$ -module consisting of all formal F -linear combinations of α -tabloids. Then Σ_n acts on α -tabloids in the obvious way: $\{t\}\pi := \{t\pi\}$. James shows that the corresponding Σ_n -module M^α is isomorphic to the permutation module $F_{R_t} \uparrow^{\Sigma_n}$.

The following element of M^α is called a *polytabloid*:

$$e_t := \sum_{\pi \in C_t} \text{sgn } \pi \{t\pi\}.$$

The subspace of M^α spanned by polytabloids is a Σ_n -module called the Specht module S^α . For compactness we use $X^- := \sum_{\sigma \in X} \text{sgn}(\sigma)\sigma$, whenever $X \subseteq \Sigma_n$. So in this notation $e_t = \{t\}C_t^-$.

The following important combinatorial result is one half of [4, 13.5].

Lemma 2.1. *Suppose that T is an α -tableau that contains repeated entries in one of its columns. Then $TC_t^- = 0$.*

Proof. By hypothesis there exist integers r, s, c , with $r \neq s$, such that $T_{r,c} = T_{s,c}$. Then $\pi := (t_{r,c}, t_{s,c})$ is a transposition in C_t . So $\pi C_t^- = \text{sgn}(\pi) C_t^- = -C_t^-$. But $T\pi = T$. It follows that $TC_t^- = T\pi C_t^- = -TC_t^-$. As $\text{char}(F) \neq 2$, this implies that $TC_t^- = 0$. \square

3. BRANCHING RULES FOR ENDOMORPHISM RINGS

As in [3], we use E_n denote the sum, in $F\Sigma_n$, of all transpositions in Σ_n . Each $F\Sigma_n$ -module M comes equipped with a ring homomorphism $F\Sigma_n \rightarrow \text{End}_F(M)$.

Theorem 3.1. *Let λ be a partition of n . Then the image of E_{n+1} generates $\text{End}_{F\Sigma_{n+1}}(S^\lambda \uparrow^{\Sigma_{n+1}})$ and the image of E_{n-1} generates $\text{End}_{F\Sigma_{n-1}}(S^\lambda \downarrow_{\Sigma_{n-1}})$.*

Proof. Let M be the a summand of the restricted or induced module belonging to a single block of the group algebra. We change notation from $n+1$ to n in the first case, and from $n-1$ to n in the second case, so that M is an $F\Sigma_n$ -module, and E_n plays the role of E_{n+1} or of E_{n-1} . Let ϵ denote the image of E_n in $\text{End}_F(M)$. Then [4] and the authors' paper [3] shows that M has the following properties:

- (1) M has a Specht series

$$0 = M_0 \subset M_1 \subset \dots \subset M_m = M,$$

such that $M_u/M_{u-1} \cong S^{\lambda_u}$ where λ_u is a partition of n . Moreover, λ_v is a one-box-shift of λ_u , for each $v \neq u$.

- (2) The labelling partitions satisfy $\lambda_1 \triangleleft \dots \triangleleft \lambda_m$.
 (3) The endomorphism ϵ has minimal polynomial $(x-r)^m$. Here ϵ acts as a fixed scalar r on each quotient module M_u/M_{u-1} , for $u = 1, \dots, m$.

It is known that the scalar r is the sum, taken modulo p , of the residues of the nodes of the Young diagram of any partition λ_u that is associated with the given p -block of Σ_n . For convenience, we define $M_u := 0$, when $u \leq 0$.

Let θ be any $F\Sigma_n$ -homomorphism of M . We shall prove by induction that given $i \geq 0$ there exists a polynomial $f_i \in F[x]$ such that $M_u(\theta - f_i(\epsilon)) \subseteq M_{u-i}$, for $u = 1, \dots, m$. So $\theta = f_m(\epsilon)$ lies in the subalgebra generated over F by ϵ and 1_M , thus proving the Theorem.

As $p \neq 2$ and $\lambda_u \triangleleft \lambda_v$ for $u < v$, it follows from Corollary 13.17 of [4] that there are no non-zero homomorphisms $M_u/M_{u-1} \rightarrow M_v/M_{v-1}$, when $u < v$. This implies in particular that $M_u\theta \subseteq M_u$, for each $u = 1, \dots, m$ (see Lemma 3.2 of [3] for details). This proves the base case for our induction.

Suppose then that $i > 0$ is an integer such that there exists a polynomial $f_i \in F[x]$ such that $M_u(\theta - f_i(\epsilon)) \subseteq M_{u-i}$, for $u = 1, \dots, m$. Then there exist well-defined Σ_n -homomorphisms $\psi_{u,i} : M_u/M_{u-1} \rightarrow M_{u-i}/M_{u-i-1}$ given by $(v + M_{u-1})\psi_{u,i} = v(\theta - f_i(\epsilon)) + M_{u-1}$, for all $v \in M_u$.

Now $M_u(\epsilon - r)^i \subseteq M_{u-i}$, for all u . As ϵ has minimal polynomial of degree m , the induced maps $(\epsilon - r)_u^i : M_u/M_{u-1} \rightarrow M_{u-i}/M_{u-i}$ are non-zero, for $u > i$. But λ_{u-i} is a one-box-shift of λ_u . So $\text{Hom}_{F\Sigma_n}(\lambda_u, \lambda_{u-i})$ is a 1-dimensional F -space, according to Corollary 6.7. We deduce that $(\epsilon - r)_u^i$ is a basis for the space $\text{Hom}_{F\Sigma_n}(\lambda_u, \lambda_{u-i})$, for all u, i .

The previous two paragraphs imply that there exist $\Lambda_u \in F$ such that $\psi_{u,i} = \Lambda_u(\epsilon - r)^i$, for $u = 1, \dots, m$. Then $M_u(\theta - f_{i+1}(\epsilon)) \subseteq M_{u-i}$, for $u = 1, \dots, m$, where $f_{i+1}(x) = f_i(x) + \Lambda_m(x - r)^i$, if we can show that all Λ_u are equal to Λ_m . Lemma 3.1 of [3] furnishes us with an element $\tau \in M$ such that $\tau_u := \tau \prod_{i=u+1}^m (\epsilon - r)$ belongs to $M_u \setminus M_{u-1}$, for each $u \in \{1, \dots, m\}$. Thus

$$\begin{aligned} \Lambda_u \tau_{u-i}(\epsilon - r)^i &= \tau_u(\theta - f_i(\epsilon)), \quad \text{mod } M_{u-i-1}, \\ &= \tau(\epsilon - r)^{m-u}(\theta - f_i(\epsilon)) \\ &= \tau(\theta - f_i(\epsilon))(\epsilon - r)^{m-u}, \quad \text{as } \theta \in \text{End}_{F\Sigma_n}(M), \\ &\equiv \Lambda_m \tau(\epsilon - r)^i(\epsilon - r)^{m-u}, \quad \text{mod } M_{m-i-1}, \\ &\equiv \Lambda_m \tau(\epsilon - r)^i, \quad \text{mod } M_{u-i-1}. \end{aligned}$$

We conclude that $\Lambda_u = \Lambda_m$. This completes the proof of the inductive step, and hence of the Corollary.

Now consider the map from the polynomial algebra $F[x]$ to the summand associated to a single block of the endomorphism algebra that sends x to multiplication by E_n . We have shown that this map is surjective and has kernel generated by a power of $x - r$, where E_n acts as the scalar r on each simple module in the block. If we consider a different block, the scalar r will be different. The theorem follows. \square

Proof of Theorem 1.3. Each p -block of Σ_n has an associated positive integer w known as its weight and a p -core κ . Here κ is a partition of $n - pw$ that has no hook-lengths divisible by p . A Specht module S^μ belongs to the block if and only if the p -core of μ coincides with κ . It is well known that the p -core of a partition is determined by the multi-set of residues (mod p) of the nodes in its Young diagram. Fix the residue r of a removable node in $[\lambda]$. Then all partitions that can be obtained by removing a single node of p -residue r from λ belong to the same p -block, call it b , of Σ_n .

The main result of [3], Theorem 3.4, shows that $S^\lambda \downarrow_{\Sigma_n} b$ is indecomposable. As this is an ideal and a direct summand of $\text{End}_{F\Sigma_n}(S^\lambda \downarrow_{\Sigma_n})$, it is a block of this algebra. Theorem 3.1 implies that $\text{End}_{F\Sigma_n}(S^\lambda \downarrow_{\Sigma_n})b$ is generated by over F by the image of E_n . Let m be the number of removable nodes in $[\lambda]$ that have p -residue r . Then E_n has minimal polynomial $(x - r)^m$. It follows that $\text{End}_{F\Sigma_n}(S^\lambda \downarrow_{\Sigma_n})b$ has the structure of the truncated polynomial ring $F[x]/(x^m)$. \square

The proof of Theorem 1.4 is almost identical, and is omitted.

Finally we note the following fact about Carter-Payne homomorphisms between Specht modules. The proof is omitted, although it can easily be demonstrated by applying the methods used in the proof of Theorem 3.1.

Proposition 3.2. *Suppose that λ is a partition of $n + 1$ and that r is the p -residue of a removable node in $[\lambda]$. Let $\lambda_1 \triangleleft \dots \triangleleft \lambda_m$ be all the partitions of n whose Young diagram can be obtained by removing a single node of p -residue r from $[\lambda]$. Let f_i be a non-zero $F\Sigma_n$ -homomorphism $S^{\lambda_i} \rightarrow S^{\lambda_{i-1}}$, for each $i = 2, \dots, m$. Then for $1 \leq j < i \leq m$, the composite homomorphism $f_i \circ f_{i-1} \circ \dots \circ f_{j+2} \circ f_{j+1}$ spans $\text{Hom}_{F\Sigma_n}(S^{\lambda_i}, S^{\lambda_j})$. In particular, all these homomorphisms are non-zero.*

4. SEMI-STANDARD HOMOMORPHISMS

We now discuss homomorphisms between Specht and permutation modules. If β is a partition of n , it is useful to describe the permutation module M^β using

certain α -tableaux. Following James, an α -tableau T is of type β if it has β_i entries equal to i , for each $i \geq 1$. The action of Σ_n on α -tableau restricts to an action on the α -tableau of type β . Under this action, the F -span of the α -tableau of type β form a permutation module that is isomorphic to M^β .

Let S and T be α -tableau of type β . We write

$$S \approx T, \quad \text{if } S = T\pi, \text{ for some } \pi \in R_t.$$

Then \approx is an equivalence relation of α -tableau. We also say that S is *row equivalent* to T . James defines the Σ_n homomorphism $\theta_T : M^\alpha \rightarrow M^\beta$ by $\{t\} \theta_T := \sum_{S \approx T} S$.

A tableau T is said to be *semi-standard* if the numbers are nondecreasing along the rows of T and strictly increasing down the columns of T . Let $\widehat{\theta}_T$ denote the restriction of θ_T to $S^\alpha \subseteq M^\alpha$. James shows in [4, 13.13] that the set

$$(3) \quad \{\widehat{\theta}_T \mid T \text{ a semi-standard } \alpha\text{-tableau of type } \beta\}$$

forms a basis for $\text{Hom}_{F\Sigma_n}(S^\alpha, M^\beta)$.

A useful result of James identifies S^β as an intersection of the kernels of certain Σ_n -homomorphisms on M^β . Let β be a partition of n that has k parts. Then [4, 17.18] shows that

$$(4) \quad S^\beta := \bigcap_{i=1}^{k-1} \bigcap_{r=0}^{\beta_{i+1}-1} \ker \psi_{i,r}.$$

Here for each pair i, r , let $\nu := (\beta_1, \beta_2, \dots, \beta_{i-1}, \beta_i + \beta_{i+1} - r, r, \beta_{i+2}, \dots)$. Then $\psi_{i,r} : M^\beta \rightarrow M^\nu$ is defined by setting $\{t\} \psi_{i,r}$ as the sum of all ν -tabloids $\{s\}$ such that row $(i+1)$ of $\{s\}$ is a subset of size r of row $(i+1)$ of $\{t\}$.

The action of $\psi_{i,r}$ on α -tableaux of type β is particularly easy to describe. If T is one such tableau then $T \psi_{i,r}$ is the sum of all α -tableaux that can be obtained from T by changing all but r symbols $i+1$ to i . For example consider the effect of $\psi_{1,2}$ on a $(4, 3)$ -tableau of type $(3^2, 1)$:

$$\begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} \xrightarrow{\psi_{1,2}} \begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & 3 \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 1 & 2 & 3 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 1 & 3 & \\ \hline \end{array}.$$

In terms of semistandard homomorphisms, this gives

$$\theta_{\begin{array}{|c|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & \\ \hline \end{array}} \psi_{1,2} = 4 \theta_{\begin{array}{|c|c|c|} \hline 1 & 1 & 1 \\ \hline 2 & 2 & 3 \\ \hline \end{array}} + \theta_{\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 1 & 2 & 3 & \\ \hline \end{array}}.$$

It is convenient to use the following concise notation

$$\psi_i := \psi_{i, \beta_{i+1}-1}.$$

In particular $T \psi_i$ is the sum of all α -tableaux of that can be obtained from T by changing one symbol $i+1$ to an i .

For the rest of this paper let $a < b$ be integers such that $\alpha_a > \alpha_{a+1}$ and $\alpha_{b-1} > \alpha_b$. So (a, α_a) is a removable node of α and $(b, \alpha_b + 1)$ is an addable node of α . We fix the partition β of n defined by

$$(5) \quad \beta_i = \begin{cases} \alpha_a - 1, & \text{if } i = a; \\ \alpha_b + 1, & \text{if } i = b; \\ \alpha_i, & \text{if } i \neq a, b. \end{cases}$$

We say that β is a *one-box-shift* of α . Note that $\alpha \triangleright \beta$ in the dominance order.

Our first task is to enumerate all semi-standard α -tableau of type β . So let T be such a tableau. We claim that

$$(6) \quad T(i, j) = i, \quad \text{for all } (i, j) \in [\alpha], \text{ unless } a \leq j < b \text{ and } j = \alpha_i.$$

For, $T(i, j) \geq i$, as T is column strict. So all symbols i occur in the top i rows of T . In particular, the symbols $1, \dots, a-1$ occupy all entries in rows $1, \dots, a-1$.

Now suppose that $a \leq i < b$. We prove by induction on i that $T(i, j) = i$, for $1 \leq j < \alpha_i$. The base case $i = a$ holds because the $\alpha_a - 1$ symbols in T must occupy all but the last entry in row a of T (there is no room for them further up). Now let $a < i$. Then there are $-1 + \sum_{u=1}^{i-1} \alpha_u$ symbols in T equal to one of $1, \dots, i-1$. These occupy all but one of the $\sum_{u=1}^{i-1} \alpha_u$ nodes in the first $i-1$ rows of T . It follows that at most one symbol i in T does not belong to row i . This proves the inductive step.

Finally, suppose that $b \leq i$. Then there are $\sum_{u=1}^i \alpha_u$ symbols in T equal to one of $1, \dots, i$ and $\sum_{u=1}^{i-1} \alpha_u$ nodes in the first i rows of T . It follows from this that every entry in the i -th row of T is equal to i . This completes the proof of (6).

Next we define

$$\hat{T}(u) := T(u, \alpha_u), \quad \text{for } u = a, \dots, b-1.$$

Suppose that $a < i \leq b$. Then there are β_i -symbols in T equal to i . At most one of them does not belong to the i -th row of T . Moreover, in view of (6), if a symbol i does not belong to row i , it occupies the end of a row between a and $i-1$ inclusive. This shows that there exists a unique u with $a \leq u \leq i$ such that $\hat{T}(u) = i$. It follows that $\hat{T} : \{a, \dots, b-1\} \rightarrow \{a+1, \dots, b\}$ is a bijection. Moreover, $\hat{T}(i) \geq i$, and $\hat{T}(i+1) > \hat{T}(i)$, if $\alpha_{i+1} = \alpha_i$. We call any such bijection \hat{T} the *semi-standard α -bijection of type β* associated with T .

We can recover the tableau T from the bijection \hat{T} , using property (6).

Let $u_1 := a$ and inductively define $u_i := \hat{T}(u_{i-1})$, for $i = 2, 3, \dots$, if $u_{i-1} < b$. This gives a finite set of cardinality $m \geq 1$:

$$\{T\} := \{u_1 = a < u_2 < \dots < u_m\}.$$

We claim that $\hat{T}(u) \neq u$ if and only if $u \in \{T\}$. We prove this by induction on $u \in \{a, \dots, b-1\}$. For, suppose that $\hat{T}(u) \neq u$. Then there exists v with $a \leq v < u$ and $\hat{T}(v) = u$. The inductive hypothesis implies that $v \in \{T\}$. So $v = u_i$, for some $i \geq 1$, whence $u = \hat{T}(u_i) = u_{i+1}$ also belongs to $\{T\}$. Note that $b = u_{m+1}$. If we temporarily extend \hat{T} to a permutation of the set $\{a, \dots, b\}$ via $\hat{T}(b) := a$, then this shows that \hat{T} is the cycle permutation $(a = i_1, i_2, \dots, i_m, i_{m+1} = b)$. So we can recover \hat{T} from $\{T\}$.

In order for a subset X of $\{a, \dots, b-1\}$ to equal $\{T\}$ for some semi-standard α -tableau of type β , it is necessary and sufficient that $a \in X$, and if $u \in X$ and $\alpha_u = \alpha_{u+1}$, then $u+1 \in X$. We call any such X a semi-standard α -set of type β . We call $\{T\}$ the semi-standard α -set of type β associated with T (or with \hat{T}). We summarize the above discussion with:

Lemma 4.1. *The associations $T \longleftrightarrow \hat{T} \longleftrightarrow \{T\}$ establish mutually inverse bijections between the semi-standard α -tableau of type β , the semi-standard α -bijections of type β , and the semi-standard α -sets of type β .*

Given $T \leftrightarrow \hat{T} \leftrightarrow \{T\}$, we can use $\hat{\theta}_{\hat{T}}$ or $\hat{\theta}_{\{T\}}$ to denote $\hat{\theta}_T$. Moreover, we use $|T|$ or $|\hat{T}|$ to denote the cardinality of $\{T\}$.

If α has m_i parts of length i , then there are $\prod_{\alpha_a < i < \alpha_b} (m_i + 1)$ semistandard α -tableau of type β . As an example let $\alpha = (4, 3^2, 2, 1)$ and $\beta = (3^3, 2, 1^2)$. So $a = 1, b = 6$ and $m_3 = 2, m_2 = 1, m_1 = 1$. Then there are $12 = (3 + 1)(1 + 1)(1 + 1)$ semistandard α -tableau of type β . We list them along with the semistandard α -sets of type β . We use \cdot to indicate a symbol i in row i :

$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 6 \\ \hline \cdot & \cdot & 2 & \\ \hline \cdot & \cdot & 3 & \\ \hline \cdot & 4 & & \\ \hline 5 & & & \\ \hline \end{array}$ $\{1\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 5 \\ \hline \cdot & \cdot & 2 & \\ \hline \cdot & \cdot & 3 & \\ \hline \cdot & 4 & & \\ \hline 6 & & & \\ \hline \end{array}$ $\{1, 5\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 4 \\ \hline \cdot & \cdot & 2 & \\ \hline \cdot & \cdot & 3 & \\ \hline \cdot & 6 & & \\ \hline 5 & & & \\ \hline \end{array}$ $\{1, 4\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 4 \\ \hline \cdot & \cdot & 2 & \\ \hline \cdot & \cdot & 3 & \\ \hline \cdot & 5 & & \\ \hline 6 & & & \\ \hline \end{array}$ $\{1, 4, 5\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 3 \\ \hline \cdot & \cdot & 2 & \\ \hline \cdot & \cdot & 6 & \\ \hline \cdot & 4 & & \\ \hline 5 & & & \\ \hline \end{array}$ $\{1, 3\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 3 \\ \hline \cdot & \cdot & 2 & \\ \hline \cdot & \cdot & 5 & \\ \hline \cdot & 4 & & \\ \hline 6 & & & \\ \hline \end{array}$ $\{1, 3, 5\}$
$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 3 \\ \hline \cdot & \cdot & 2 & \\ \hline \cdot & \cdot & 4 & \\ \hline \cdot & 6 & & \\ \hline 5 & & & \\ \hline \end{array}$ $\{1, 3, 4\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 3 \\ \hline \cdot & \cdot & 2 & \\ \hline \cdot & \cdot & 4 & \\ \hline \cdot & 5 & & \\ \hline 6 & & & \\ \hline \end{array}$ $\{1, 3, 4, 5\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 2 \\ \hline \cdot & \cdot & 3 & \\ \hline \cdot & \cdot & 6 & \\ \hline \cdot & 4 & & \\ \hline 5 & & & \\ \hline \end{array}$ $\{1, 2, 3\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 2 \\ \hline \cdot & \cdot & 3 & \\ \hline \cdot & \cdot & 5 & \\ \hline \cdot & 4 & & \\ \hline 6 & & & \\ \hline \end{array}$ $\{1, 2, 3, 5\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 2 \\ \hline \cdot & \cdot & 3 & \\ \hline \cdot & \cdot & 4 & \\ \hline \cdot & 6 & & \\ \hline 5 & & & \\ \hline \end{array}$ $\{1, 2, 3, 4\}$	$\begin{array}{ c c c c } \hline \cdot & \cdot & \cdot & 2 \\ \hline \cdot & \cdot & 3 & \\ \hline \cdot & \cdot & 4 & \\ \hline \cdot & 5 & & \\ \hline 6 & & & \\ \hline \end{array}$ $\{1, 2, 3, 4, 5\}$

For the rest of the paper we let

$$\{R\} := \{i \mid (i, \alpha_i) \text{ is a removable node of } [\alpha], a \leq i \leq b - 1\}.$$

The associated semi-standard α -bijection of type β is denoted by \hat{R} , and the tableau by R . A subset of $\{R\}$ that contains a is called a *removable α -set of type β* , or simply a *removable set*. Clearly each removable α -set of type β is also a semi-standard α -set of type β . Each semi-standard set $\{T\}$ contains a largest removable set $\{N\} = \{T\} \cap \{R\}$.

We say that a pair of removable nodes are *adjacent* if there is no removable node of $[\alpha]$ between them. Likewise, if $i, j \in \{R\}$, we say that i and j are adjacent if the removable nodes (i, α_i) and (j, α_j) are adjacent; if $i < j$ this means that $\alpha_{i+1} = \alpha_j$.

5. RELATIONS BETWEEN SEMI-STANDARD HOMOMORPHISMS

Throughout this section T is a fixed semi-standard α -tableau of type β .

The hooks-lengths in the $(\alpha_b + 1)$ -th column of $[\alpha]$ are h_1, \dots, h_{b-1} , where

$$h_i := \alpha_i - \alpha_b + b - i - 1, \quad \text{for } 1 \leq i \leq b - 1.$$

For convenience we set $h_i := 0$, for each $i \geq b$. Note that

$$(7) \quad \begin{aligned} h_i - h_j &= (\alpha_i - \alpha_j) + (j - i), & \text{for } 1 \leq i, j \leq b - 1 \text{ and} \\ h_{b-1} - h_b &= \alpha_{b-1} - \alpha_b. \end{aligned}$$

We begin with a useful technical result.

Lemma 5.1. *Let $U \approx T$, let $i \geq 0$ and let V be an α -tableau that is obtained from U by changing a set of $r \geq 1$ symbols $i + 1$ to i , with at least one of the changes occurring in row $(i + 1)$ of U . Suppose that $UC_t^- \neq 0$ and that $VC_t^- \neq 0$. Then $r = 1$ and $\hat{T}(i) > i + 1$ and $\hat{T}(i + 1) = i + 1$.*

Proof. Let $c \geq 1$ be such that $U(i + 1, c) = i + 1$ and $V(i + 1, c) = i$. As $UC_t^- \neq 0$ and $VC_t^- \neq 0$, separate applications of Lemma 2.1 show that $U(i, c) \neq i + 1$ and $V(i, c) \neq i$. Thus $U(i, c) \neq i, i + 1$. As $U \approx T$, we deduce that $\hat{T}(i) > i + 1$. It follows from this that $\hat{T}(i + 1) = i + 1$. In particular row $i + 1$ contains all α_{i+1} -symbols $i + 1$ in U . Now (i, c) is the unique node in row i of U that does not contain i . We conclude that $r = 1$. \square

We will repeatedly, and without further comment, use the fact that S^α is generated by e_t as a Σ_n -module. Moreover for all i and r

$$(8) \quad e_t \widehat{\theta}_T \psi_{i,r} = \sum_{U \approx T} U \psi_{i,r} C_t^-,$$

as both $\widehat{\theta}_T$ and $\psi_{i,r}$ are Σ_n -homomorphisms.

Lemma 5.2. *Let $i \geq 1$ and $1 \leq r \leq \beta_{i+1} - 1$ be such that $\widehat{\theta}_T \psi_{i,r} \neq 0$. Then $a \leq i < b$ and $r = \beta_{i+1} - 1$.*

Proof. Let $i \geq 0$, let $U \approx T$ and let V be a tableau occurring in $U \psi_{i,r}$.

Suppose that $1 \leq r < \beta_{i+1} - 1$. As T is semi-standard, Lemma 4.1 implies that at most one symbol $i + 1$ in U does not belong to row $(i + 1)$. But V is the result of changing $\beta_{i+1} - r \geq 2$ symbols in U from $i + 1$ to i . So one of the changes occurs in row $(i + 1)$ of U . It then follows from Lemma 5.1 that either $U C_t^- = 0$ or $V C_t^- = 0$.

Suppose that $i < a$ or $b < i$. Then all symbols $i + 1$ in U belong to row $(i + 1)$ of U , and Lemma 5.1 implies that $U C_t^- = 0$ or $V C_t^- = 0$.

In all cases we have $U C_t^- \psi_i = U \psi_i C_t^- = \sum V C_t^- = 0$. We then use (8) to conclude that $\widehat{\theta}_T = 0$, which is contrary to hypothesis. \square

Lemmas 5.3, 5.4 and 5.5 will show that the image of $\widehat{\theta}_T$ under ψ_i is either zero or a specified non-zero integer multiple of a semi-standard homomorphism from S^α into the permutation module M^γ . Here γ is the composition

$$\gamma_j = \begin{cases} \beta_i + 1, & \text{if } j = i; \\ \beta_{i+1} - 1, & \text{if } j = i + 1; \\ \beta_j, & \text{otherwise.} \end{cases}$$

Notice that $\beta \triangleleft \gamma \triangleleft \alpha$.

Lemma 5.3. *Let $a < i \leq b - 1$ be such that $i \notin \{T\}$. Then*

$$\widehat{\theta}_T \psi_i = \begin{cases} \widehat{\theta}_W, & \text{if } i + 1 \in \{T\}; \\ 0, & \text{otherwise.} \end{cases}$$

Here W is obtained from T by changing the single $i + 1$ in row $\widehat{T}^{-1}(i + 1)$ to an i .

Proof. Let $U \approx T$. The hypothesis on i implies that the α_i symbols i occupy the α_i columns of row i in U . So each symbol $i + 1$ which belongs to row $i + 1$ of U shares its column with a symbol i in row i of U . Let $a \leq j < b$ be such that $\widehat{T}(j) = i + 1$.

Suppose first that $j = i + 1$. Then all symbols $i + 1$ in U occur in row $(i + 1)$. It then follows from Lemma 5.1 and the first paragraph that $U \psi_i C_t^- = 0$. We conclude from (8) that $\widehat{\theta}_T = 0$ in this case.

From now on assume that $j \neq i + 1$, whence $j < i$. Then exactly one of the symbols $i + 1$ in U occurs in row j , and all the other symbols $i + 1$ belong to row $(i + 1)$ of U . It then follows from Lemma 5.1 and the first paragraph that either $U C_t^- = 0$ or $U \psi_i C_t^- = V C_t^-$, where V is the tableau obtained from U by changing the symbol $i + 1$ in row j to an i . It is obvious that $V \approx W$ and also that U is the only tableau that is row equivalent to T such that $U \psi_i C_t^- = V C_t^-$.

Conversely begin with $V \approx W$. Then V has a unique i in row j . Changing this i to an $i + 1$ produces a tableau U that is row equivalent to T . Moreover clearly $U \psi_i C_t^- = V C_t^-$.

These arguments and (8) show that $e_t \widehat{\theta}_T \psi_i = \sum_{V \approx W} V C_t^- = e_t \widehat{\theta}_W$. We conclude that in this case $\widehat{\theta}_T \psi_i = \widehat{\theta}_W$. This completes the proof. \square

Lemma 5.4. *Let $a < i \leq b - 1$ be such that $\alpha_{\widehat{T}^{-1}(i)} = \alpha_i$. Then*

$$\widehat{\theta}_T \psi_i = 0.$$

Proof. The hypothesis implies that $\widehat{T}(i-1) = i$ and $\alpha_{i-1} = \alpha_i$. Let $U \approx T$ be such that $U C_t^- \neq 0$. Then one symbol i belongs to row $(i-1)$ of U , and the remaining $\alpha_i - 1$ symbols i belong to row i of U . It then follows from Lemma 2.1 that each of the first α_i columns of U contains exactly one symbol i .

Now either $\widehat{T}(i) = i + 1$ or $\widehat{T}(i + 1) = i + 1$. In either case all symbols $i + 1$ belongs to the first α_i columns of U . It then follows from the previous paragraph that each symbol $i + 1$ in U belongs to a column that also contains a symbol i . We deduce from Lemma 2.1 that $U \psi_i C_t^- = 0$. This contradiction means that our supposition that U exists is false.

Using (8), we see that $e_t \widehat{\theta}_T \psi_i = 0$. We conclude that $\widehat{\theta}_T \psi_i = 0$. \square

Suppose that $a \leq i \leq b - 1$ is such that $\widehat{T}(i) \neq i$. Let $k \geq 1$ be such that $(i + k, \alpha_{i+k})$ is a removable node of α , and $\alpha_{i+1} = \alpha_{i+k}$. If $\widehat{T}(i) = i + 1$, set $T \vee i := T$. Otherwise, set $\{T \vee i\} := \{T\} \cup \{i + 1, \dots, i + k\}$. Then $\{T \vee i\}$ is the smallest semi-standard α -set of type β that contains $i + 1$ and $\{T\}$. There is a corresponding α -tableau $T \vee i$ and α -bijection $\widehat{T \vee i}$:

$$(9) \quad \widehat{T \vee i}(r) = \begin{cases} r + 1, & \text{if } i \leq r < i + k < b; \\ \widehat{T}(i), & \text{if } r = i + k \text{ and } \widehat{T}(i + k) = i + k; \\ \widehat{T}(r), & \text{otherwise.} \end{cases}$$

Suppose that T is such that $\widehat{T}(i+k) = i+k$. In particular (i, α_i) is a removable node of α . The following describe the semi-standard α -sets $\{S\}$ such that $S \vee i = T \vee i$:

$$\begin{array}{l} \{T\} \\ \{T\} \cup \{i + k\} \\ \{T\} \cup \{i + k - 1, i + k\} \\ \vdots \\ \{T\} \cup \{i + 2, \dots, i + k - 1, i + k\} \\ \{T\} \cup \{i + 1, i + 2, \dots, i + k - 1, i + k\} = \{T \vee i\} \end{array}$$

Lemma 5.5. *Let $a \leq i \leq b - 1$ be such that $i = a$ or $\alpha_{\widehat{T}^{-1}(i)} \neq \alpha_i$. Then*

$$\widehat{\theta}_T \psi_i = \begin{cases} (h_i - h_{i+1}) \widehat{\theta}_W, & \text{if } \widehat{T}(i) = i + 1; \\ (-1)^{|\{T \vee i\} \setminus \{T\}|} \widehat{\theta}_W, & \text{otherwise.} \end{cases}$$

Here W is obtained from $T \vee i$ by changing the unique symbol $i + 1$ in row i to an i .

Proof. Let $U \approx T$ be such that $U \psi_i C_t^- \neq 0$.

We consider first the case that $\widehat{T}(i) = i + 1$, or equivalently $T = T \vee i$. In particular there exists a column number c so that node (i, c) of U contains the symbol $i + 1$, and all other symbols $i + 1$ belong to row $(i + 1)$ of U . Lemma 5.1 then implies that $U \psi_i C_t^- = V C_t^-$, where V is obtained from U by changing the $i + 1$ in node (i, c) to an i . In particular $V \approx W$.

Lemma 2.1 implies that column c of U does not contain two symbols $i + 1$. So either (i, c) is to the right of all nodes in row $i + 1$ i.e. $c \in \{\alpha_{i+1} + 1, \dots, \alpha_i\}$, or $i + 1 \neq b$ and $(i + 1, c)$ is the unique node in row $i + 1$ of U that does not contain $i + 1$. This argument and (7) shows that there are $\alpha_i - \alpha_{i+1} + 1 = h_i - h_{i+1}$ choices for a tableau U such that $U\psi_i C_t^- = V C_t^-$. It then follows from (8) that

$$e_t \widehat{\theta}_T \psi_i = (h_i - h_{i+1}) \sum_{V \approx W} V C_t^- = (h_i - h_{i+1}) e_t \widehat{\theta}_W.$$

We conclude that in this case $\widehat{\theta}_T \psi_i = (h_i - h_{i+1}) \widehat{\theta}_W$.

Now consider the case that $\widehat{T}(i) \neq i + 1$. Then $\widehat{T}(i + 1) = i + 1$. As $\widehat{T}(i) \neq i$, this ensures that (i, α_i) is a removable node of α . Set $j := \widehat{T}(i)$. Lemma 4.1 shows that there exists $c \geq 1$ such that node (i, c) of U contains the symbol $i + j > i + 1$, and all other nodes in row i contain the symbol i . Furthermore, all symbols $i + 1$ in U occur in row $(i + 1)$ of U . Then Lemma 2.1 implies that $c \leq \alpha_{i+1}$ and $U\psi_i C_t^- = V C_t^-$, where V is obtained from U by changing the $i + 1$ in node $(i + 1, c)$ of U to an i .

Let $k \geq 1$ be such that $\alpha_{i+k} = \alpha_{i+1}$ and $(i + k, \alpha_{i+k})$ is a removable node of α . Let π be the permutation in Σ_n that cycles the entries in the nodes of U as follows:

$$\begin{aligned} \text{if } j > k \text{ then } \pi : & (i + k, c) \rightarrow \dots \rightarrow (i + 1, c) \rightarrow (i, c) \rightarrow (i + k, c); \\ \text{if } j \leq k \text{ then } \pi : & (i + j - 1, c) \rightarrow \dots \rightarrow (i + 1, c) \rightarrow (i, c) \rightarrow (i + j - 1, c). \end{aligned}$$

Then π belongs to C_t . So $\pi C_t^- = \text{sgn}(\pi) C_t^-$. Also $V\pi \approx W$ is strictly increasing down columns, and we obtain each α -tableau that is row equivalent to W as an $V\pi$ exactly once, by an appropriate choice of $U \approx T$.

Note that

$$\{T \vee i\} = \begin{cases} \{T\} \dot{\cup} \{i + 1, \dots, i + k\}, & \text{if } j > k, \\ \{T\} \dot{\cup} \{i + 1, \dots, i + j - 1\}, & \text{if } j \leq k. \end{cases}$$

So $\text{sgn}(\pi) = (-1)^{|\{T \vee i\} \setminus \{T\}|}$ does not depend on the choice of $U \approx T$. Thus

$$e_t \widehat{\theta}_T \psi_i = \sum_{U \approx T} U \psi_i C_t^- = (-1)^{|\{T \vee i\} \setminus \{T\}|} \sum_{V \approx W} V C_t^- = (-1)^{|\{T \vee i\} \setminus \{T\}|} e_t \widehat{\theta}_W.$$

It follows that in this case $\widehat{\theta}_T \psi_i = (-1)^{|\{T \vee i\} \setminus \{T\}|} \widehat{\theta}_W$. \square

6. CARTER-PAYNE HOMOMORPHISMS

We now fix a Σ_n -homomorphism $\hat{\theta} : S^\alpha \rightarrow S^\beta$. By (3) we may write

$$(10) \quad \hat{\theta} = \sum_T \Lambda_T \hat{\theta}_T, \quad \text{for certain } \Lambda_T \in F,$$

where T ranges over the semi-standard α -tableaux of type β . We aim to show that, up to a scalar multiple, there is at most one such homomorphism $\hat{\theta}$. Since the image of $\hat{\theta}$ is a submodule of S^β , it follows that $\hat{\theta} \psi_i = 0$ for all i . Fix an i . Lemmas 5.3, 5.4 and 5.5 show that for all T , $\hat{\theta}_T \psi_i$ is equal to $a_{T,i} \hat{\theta}_{S_{T,i}}$, where $a_{T,i}$ is an integer and $S_{T,i}$ is a semistandard tableau depending on T and i . Maps associated to semistandard tableaux are linearly independent. Thus the equation $\hat{\theta} \psi_i = 0$ produces several equations that the coefficients Λ_T must satisfy; for any semistandard tableau W , we have $\sum \Lambda_T a_{T,i} = 0$, where the sum is over all T such that $S_{T,i} = W$. We shall collect enough of these equations to show that the space of all Σ_n -homomorphisms from S^α to S^β is at most one-dimensional.

Lemma 6.1. *Let $\{T\}$ be a semi-standard α -set of type β . Then*

$$(-1)^{|\{T\}|} \Lambda_{\{T\}} = (-1)^{|\{T\} \cap \{R\}|} \Lambda_{\{T\} \cap \{R\}}.$$

Proof. The result is trivial if $\{T\} \subseteq \{R\}$. We prove the result by induction on $|\{T\} \setminus \{R\}|$. So we may assume that $\{T\} \setminus \{R\}$ is nonempty. In particular, we may choose $i \in \{a+1, \dots, b-1\}$ such that $\alpha_{\hat{T}^{-1}(i)} > \alpha_i$ and $\alpha_i = \alpha_{i+1}$. This forces $\hat{T}(i) = i+1$. In terms of tableaux, this can be visualized as follows. The set $\{T\}$ is associated to a series of vertical strips of entries at the right edge of the tableau, vertical strips whose lowest entry is at a removable node. We are focussing on an entry at the top of one of these strips. The inductive step consists of proving that $\Lambda_T + \Lambda_{\{T\} \setminus \{i\}} = 0$. Geometrically the end result of this lemma is to show that we need only understand the coefficients of semistandard sets associated to unions of strips of length 1 containing a single removable node.

Lemma 5.5 implies that $\widehat{\theta}_T \psi_i = \widehat{\theta}_W$, where W is the α -tableau that is obtained from T by changing the single $i+1$ in row i of T to an i .

Suppose that U is a semi-standard α -tableau of type β such that $\widehat{\theta}_U \psi_i$ is a non-zero multiple of $\widehat{\theta}_W$. Lemmas 5.3, 5.4 and 5.5 imply that either $i \notin \{U\}$ or (as $\alpha_i = \alpha_{i+1}$) both i and $i+1$ belong to $\{U\}$. Moreover $\{T\}$ and $\{U\}$ do not differ apart from in the set $\{i, i+1\}$. In the former case $\{U\} = \{T\} \setminus \{i\}$ and hence $\widehat{\theta}_U \psi_i = \widehat{\theta}_W$. In the latter case $\{U\} = \{T\}$. This completes the proof of the inductive step, and hence the proof of the lemma also. \square

From now on we concentrate on finding relations among the Λ_N , with $\{N\} \subseteq \{R\}$ a removable α -set of type β . So we consider $N \leftrightarrow \hat{N} \leftrightarrow \{N\}$ such that $\{N\}$ is an arbitrary removable α -set of type β .

Lemma 6.2. *Let $i \in \{N\}$ and $j = \hat{N}(i)$ be such that (i, α_i) and (j, α_j) are adjacent removable nodes of $[\alpha]$. Then*

$$h_i \Lambda_{\{N\}} + \Lambda_{\{N\} \setminus \{i\}} = h_j \Lambda_{\{N\}} + \Lambda_{\{N\} \setminus \{j\}}.$$

The term involving $\{N\} \setminus \{i\}$ is omitted if $i = a$, while the term involving $\{N\} \setminus \{j\}$ is omitted if $i = b-1$ (and so $j = b$; recall also that $h_b = 0$).

Proof. We assume the generic case that $i \neq a$ and $j \neq b$. The two exceptional cases follow from similar arguments.

Lemma 5.5 implies that $\widehat{\theta}_N \psi_i$ is a non-zero multiple of $\widehat{\theta}_W$. Here W is the tableau that is obtained by changing the single $i+1$ in row i of $N \vee i$ to an i . Note that $\{N \vee i\} = \{N\} \dot{\cup} \{i+1, \dots, j-1\}$. We enumerate all standard α -tableau T of type β such that $\widehat{\theta}_T \psi_i$ is a non-zero multiple of $\widehat{\theta}_W$.

Suppose that $\hat{T}(i) = i$. Then $i \neq a$. Lemma 5.3 forces $\hat{T}(i+1) \neq i+1$ and $\widehat{\theta}_T \psi_i = \widehat{\theta}_W$. Also $\{T\} = \{T \vee i\} = \{N \vee i\} \setminus \{i\}$. Thus $\{T\} \cap \{R\} = \{N\} \setminus \{i\}$ and $|\{T\} \setminus \{T\} \cap \{R\}| = j - i - 1$. Lemma 6.1 then implies that in this case

$$(11) \quad \Lambda_{\{T\}} \widehat{\theta}_T \psi_i = (-1)^{j-i-1} \Lambda_{\{N\} \setminus \{i\}} \widehat{\theta}_W.$$

From now on we assume that $\hat{T}(i) \neq i$. Then Lemma 5.5 forces $\{T \vee i\} = \{N \vee i\}$. There are three cases depending on whether $i+1, j \in \{T\}$ or not.

Suppose first that $i+1 \in \{T\}$. Then $j \in \{T\}$ and $T = T \vee i$, and Lemma 5.5 shows that $\widehat{\theta}_T \psi_i = (h_i - h_{i+1}) \widehat{\theta}_W$. But $\{T\} \cap \{R\} = \{N\}$. So $|\{T\} \setminus \{T\} \cap \{R\}| = j - i - 1$.

We conclude from Lemma 6.1 that in this case

$$(12) \quad \Lambda_{\{T\}} \widehat{\theta}_T \psi_i = (-1)^{j-i-1} (h_i - h_{i+1}) \Lambda_{\{N\}} \widehat{\theta}_W.$$

There are $j-i-1$ semi-standard sets $\{T\}$ such that $j \in \{T\}$ but $i+1 \notin \{T\}$, one for each of the rows $i+1, i+3, \dots, j$. For each such $\{T\}$ we have $\{T\} \cap \{R\} = \{N\}$. In particular $|\{T \vee i\} \setminus \{T\} \cap \{R\}| = j-i-1$. It then follows from Lemmas 5.5 and 6.1 that

$$(13) \quad \Lambda_{\{T\}} \widehat{\theta}_T \psi_i = (-1)^{j-i+1} \Lambda_{\{N\}} \widehat{\theta}_W.$$

The final possibility is that $i+1, j \notin \{T\}$. Then $\{T\} = \{N\} \setminus \{j\}$ is a removable set and $|\{T \vee i\} \setminus \{T\} \cap \{R\}| = j-i$. So by Lemmas 5.5 and 6.1 we have

$$(14) \quad \Lambda_{\{T\}} \widehat{\theta}_T \psi_i = (-1)^{j-i} \Lambda_{\{N\} \setminus \{j\}} \widehat{\theta}_W.$$

We now apply ψ_i to $\widehat{\theta}$, expand as a linear combination of semi-standard homomorphisms, and examine the coefficient of $\widehat{\theta}_W$ in this expansion. Multiplying (11), (12), (13) and (14) by $(-1)^{j-i-1}$, we get

$$\Lambda_{\{N\} \setminus \{i\}} + (h_i - h_{i+1}) \Lambda_{\{N\}} - \Lambda_{\{N\} \setminus \{j\}} + (j-i-1) \Lambda_{\{N\}} = 0.$$

Using (7) and the fact that $\alpha_{i+1} = \alpha_j$, we can rewrite this as

$$(h_i - h_j) \Lambda_{\{N\}} + \Lambda_{\{N\} \setminus \{i\}} = \Lambda_{\{N\} \setminus \{j\}}.$$

□

We now prove the main result of this section.

Theorem 6.3. $(-1)^{|N|} \Lambda_N = (-1)^{|R|} \left(\prod_{u \in \{R\} \setminus \{N\}} h_u \right) \Lambda_R.$

Proof. We shall prove this theorem by repeated applications of Lemma 6.2, with $i \neq a$. To do this, we use a partial order on removable α -tableau of type β . Suppose that $\{S\}, \{T\} \subseteq \{R\}$, with $a \in \{S\}, \{T\}$. We write $\{S\} \triangleleft \{T\}$ if $\{S\} = \{T\} \setminus \{i\}$, for some i , or if there exists (i, α_i) and (j, α_j) adjacent removable nodes of $[\alpha]$ such that $i > j$ and $\{S\} \setminus \{T\} = \{i\}$ and $\{T\} \setminus \{S\} = \{j\}$. Let $<$ be the partial order generated by \triangleleft . We shall prove the theorem by induction on $<$. The base case is $\{N\} = \{R\}$, when the conclusion is trivially true. Assume then that $\{N\} \subsetneq \{R\}$.

Suppose first that $b-1 \notin \{N\}$. We apply Lemma 6.2 to $\{N\} \cup \{b-1\}$ to get

$$\Lambda_N = (-h_{b-1}) \Lambda_{\{N\} \cup \{b-1\}}.$$

But $\{N\} \triangleleft \{N\} \cup \{b-1\}$. So the inductive hypothesis gives

$$\Lambda_{\{N\} \cup \{b-1\}} = \prod_{u \in \{R\} \setminus \{N\} \cup \{b-1\}} (-h_u) \Lambda_R.$$

The claimed expression for Λ_N follows in this case.

Suppose then that $b-1 \in \{N\}$. Then we can find adjacent removable nodes (i, α_i) and (j, α_j) of $[\alpha]$ such that $i > j > a$ and $i \in \{N\}$ but $j \notin \{N\}$. Lemma 6.2 implies that

$$\Lambda_{\{N\}} = (h_j - h_i) \Lambda_{\{N\} \cup \{j\}} + \Lambda_{\{N\} \cup \{j\} \setminus \{i\}}.$$

But $\{N\} \triangleleft \{N\} \cup \{j\}$ and $\{N\} \triangleleft \{N\} \cup \{j\} \setminus \{i\}$. So by the induction hypothesis we have

$$\begin{aligned} (h_j - h_i)\Lambda_{\{N\} \cup \{j\}} &= (-h_i + h_j) \prod_{u \in \{R\} \setminus \{N\} \cup \{j\}} (-h_u) \Lambda_R; \\ \Lambda_{\{N\} \cup \{j\} \setminus \{i\}} &= (-h_j) \prod_{u \in \{R\} \setminus \{N\} \cup \{j\}} (-h_u) \Lambda_R. \end{aligned}$$

Adding, we obtain the inductive step. \square

Our corollary is a precise form of Theorem 1.5.

Corollary 6.4. *The F -space $\text{Hom}_{F\Sigma_n}(S^\alpha, S^\beta)$ is at most one dimensional. Every non-zero Σ_n -homomorphism $S^\alpha \rightarrow S^\beta$ is a scalar multiple of*

$$\sum_T \left((-1)^{|T|} \prod_{u \in \{R\} \setminus \{T\}} h_u \right) \widehat{\theta}_T.$$

Proof. This follows at once from Lemma 6.1 and Theorem 6.3. \square

From now on we assume that

$$\widehat{\theta} = \sum_T \left((-1)^{|T|} \prod_{u \in \{R\} \setminus \{T\}} h_u \right) \widehat{\theta}_T.$$

Lemma 6.5. *Over \mathbb{Z} we have $\widehat{\theta} \psi_{i,r} = 0$, unless $i = a$ and $r = \beta_{a+1} - 1$.*

Proof. In view of Lemma 5.2, we need only show that $\widehat{\theta} \psi_i = 0$, for $i = a+1, \dots, b-1$. Fix i with $a < i < b$, and let W be an α -tableau such that there exists a semi-standard α -tableau X of type β such that $\widehat{\theta}_X \psi_i$ is a nonzero multiple of $\widehat{\theta}_W$. We enumerate all such tableau X .

Suppose first that (i, α_i) is not a removable node of $[\alpha]$. Then Lemmas 5.2, 5.3, 5.4 and 5.5 show that there is a semi-standard α -tableau T of type β with $i \notin \{T\}$ such that $X = T$ or S , where $\{S\} := \{T\} \cup \{i\}$. Moreover, Lemmas 5.3 and 5.5 and the definition of θ show that

$$(\Lambda_T \widehat{\theta}_T + \Lambda_S \widehat{\theta}_S) \psi_i = (-1)^{|T|} (1-1) \left(\prod_{u \in \{R\} \setminus \{T\}} h_u \right) \widehat{\theta}_W = 0.$$

Suppose then that (i, α_i) is a removable node of $[\alpha]$. Let $j > i$ be such that (j, α_j) is a removable node of $[\alpha]$ and $\alpha_j = \alpha_{i+1}$. So (i, α_i) and (j, α_j) are adjacent removable nodes. Then Lemmas 5.2, 5.3, 5.4 and 5.5 show that there is a semi-standard α -tableau T of type β with $i, i+1 \in \{T\}$, such that $X = T \vee i$ or $\{X\} = \{T\} \setminus \{i\}$. Thus $\{X\}$ is one of the following semi-standard α -sets:

$$\begin{aligned} &\{T\}, \\ \{S_i\} &:= \{T\} \setminus \{i\}, \\ \{S_{i+1}\} &:= \{T\} \setminus \{i+1\}, \\ \{S_{i+2}\} &:= \{T\} \setminus \{i+1, i+2\}, \\ &\vdots \\ \{S_{j-1}\} &:= \{T\} \setminus \{i+1, i+2, \dots, j-1\}, \\ \{S_j\} &:= \{T\} \setminus \{i+1, i+2, \dots, j-1, j\}. \end{aligned}$$

Lemmas 5.3 and 5.5 and show that the coefficient of $\widehat{\theta}_W$ in $\theta\psi_a$ is given by

$$(\Lambda_T \widehat{\theta}_T + \sum_{u=i}^j \Lambda_{S_u} \widehat{\theta}_{S_u}) \psi_i = (-1)^{|T|} ((h_i - h_{i+1}) - h_i + (j - i - 1) + h_j) \left(\prod_{u \in \{R\} \setminus \{T\}} h_u \right) \widehat{\theta}_W = 0,$$

as $h_j - h_{i+1} = i + 1 - j$, by (7) using the fact that $\alpha_j = \alpha_{i+1}$. \square

Let γ be the partition corresponding to the following composition of n :

$$\alpha_1, \dots, \alpha_a, \alpha_{a+1} - 1, \alpha_{a+2}, \dots, \alpha_{b-1}, \alpha_b + 1, \alpha_{b+1}, \dots$$

We now deal with the exceptional test function ψ_a .

Lemma 6.6. *Over \mathbb{Z} we have $\text{Im } \hat{\theta}\psi_a \subseteq h_a M^\gamma$.*

Proof. We retain the notation of W, T from the previous lemma, with $i = a$ and j such that (j, α_j) be the highest removable node of $[\alpha]$ below (a, α_a) . Equivalently $\alpha_{j+1} < \alpha_j$ and $\alpha_j = \alpha_{a+1}$. The semi-standard sets that contribute to $\widehat{\theta}_W$ can be enumerated as $\{T\}, \{S_{i+1}\}, \dots, \{S_j\}$ i.e. the same as in Lemma 6.5, but with $\{S_i\} = \{T\} \setminus \{a\}$ omitted (because it does not contain a , and hence is not a semi-standard α -set of type β). So the coefficient of $\widehat{\theta}_W$ in $\theta\psi_a$ is given by

$$\begin{aligned} (\Lambda_T \widehat{\theta}_T + \sum_{u=a+1}^j \Lambda_{S_u} \widehat{\theta}_{S_u}) \psi_a &= (-1)^{|T|} ((h_a - h_{a+1}) + (j - a - 1) + h_j) \left(\prod_{u \in \{R\} \setminus \{T\}} h_u \right) \widehat{\theta}_W \\ &= (-1)^{|T|} h_a \left(\prod_{u \in \{R\} \setminus \{T\}} h_u \right) \widehat{\theta}_W. \end{aligned}$$

\square

We also obtain the following criterion for the existence of a non-zero homomorphism. It is a special case of a result of Carter and Payne [2]. The ‘if’ part is also true over fields of characteristic 2.

Corollary 6.7. *Let β be a one-box-shift of α , with $\beta_a = \alpha_a - 1$ and $\beta_b = \alpha_b + 1$, for $a < b$. Then $\text{Hom}_{F\Sigma_n}(S^\alpha, S^\beta) \neq 0$ if and only if $\alpha_a - a \equiv \beta_b - b \pmod{p}$.*

Proof. Lemmas 6.5 and 6.6 show that, over the field F of characteristic p , the image of θ belongs to S^β if and only if $h_a \equiv 0 \pmod{p}$ (for the only if, for $T = R$ we have $\theta\psi_a \neq 0$ if $p \nmid h_a$ as in that case $\prod_{u \in \{R\} \setminus \{T\}} h_u = 1$). The result follows, as $h_a = (\alpha_a - \alpha_b) + (b - a - 1) = (\alpha_a - a) - (\beta_b - b)$. \square

7. JANTZEN FILTRATION

We let $\langle \cdot, \cdot \rangle$ denote the symmetric bilinear form for which the β -tabloids form an orthonormal basis of M^β . This form is Σ_n -invariant. Over \mathbb{Q} we have a direct sum decomposition:

$$M_{\mathbb{Q}}^\beta = S_{\mathbb{Q}}^\beta \oplus S_{\mathbb{Q}}^{\beta\perp},$$

where $S_{\mathbb{Q}}^{\beta\perp} = \{m \in M_{\mathbb{Q}}^\beta \mid \langle m, S_{\mathbb{Q}}^\beta \rangle = 0\}$ is the perpendicular space to $S_{\mathbb{Q}}^\beta$. Now $S_{\mathbb{Z}}^\beta \oplus S_{\mathbb{Z}}^{\beta\perp}$ is a full sublattice of $M_{\mathbb{Z}}^\beta$ (in the sense that both have the same \mathbb{Z} -rank). However the quotient lattice $M_{\mathbb{Z}}^\beta / S_{\mathbb{Z}}^\beta \oplus S_{\mathbb{Z}}^{\beta\perp}$ can be quite large. Over a field F of characteristic p , the restriction of the form $\langle \cdot, \cdot \rangle$ to S^β is generally degenerate. Over \mathbb{Z} , the module $S_{\mathbb{Z}}^\beta$ has a sequence of Jantzen submodules:

$$J^i(S_{\mathbb{Z}}^\beta) := \{x \in S_{\mathbb{Z}}^\beta \mid \langle x, S_{\mathbb{Z}}^\beta \rangle \subseteq p^i \mathbb{Z}\}.$$

The images of these modules under the decomposition map give a corresponding filtration of the Specht module $S_F^\beta = S_{\mathbb{Z}}^\beta / pS_{\mathbb{Z}}^\beta$. We note that

$$(15) \quad J^i(S_{\mathbb{Z}}^\beta) = S_{\mathbb{Z}}^\beta \cap (p^i M_{\mathbb{Z}}^\beta + S_{\mathbb{Z}}^{\beta \perp}).$$

See [7, 1.2] for example. We prove Theorem 1.6 in the following form:

Theorem 7.1. *Let α, β be partitions of n , with $\beta_a = \alpha_a - 1$ and $\beta_b = \alpha_b + 1$, for $a < b$, and $\beta_j = \alpha_j$, for $j \neq a, b$. Suppose that $p^i | (\alpha_a - \beta_b + b - a)$, for some $i \geq 0$. Then each $\hat{\theta} \in \text{Hom}_{\Sigma_n}(S^\alpha, S^\beta)$ satisfies $\text{Im}(\hat{\theta}) \subseteq J^i(S^\beta)$.*

Proof. In view of Corollary 6.4, we may assume that $\hat{\theta}$ is the map of the previous section, defined over \mathbb{Z} . Recall that the polytabloid $e_t := \{t\}C_t^-$ generates S^α . Here t is a fixed α -tableau, and C_t^- is the signed column stabilizer sum of t . Also $h_a := (\alpha_a - \beta_b + b - a)$. The strategy of this proof is very simple. In (17) below we produce an ‘error term’ $\mathcal{E}_t \in M^\beta$ and show that

$$e_t \hat{\theta} - h_a \mathcal{E}_t \in S_{\mathbb{Z}}^\beta.$$

Now over \mathbb{Z} , the image of $\hat{\theta}$ is contained in $S_{\mathbb{Z}}^{\beta \perp}$. So by (15), $e_t \hat{\theta} - h_a \mathcal{E}_t$ belongs to $J^i(S_{\mathbb{Z}}^\beta)$, which is what we need to prove.

If T is a semistandard α -tableau of type β then there is a corresponding semi-standard homomorphism $\widehat{\theta}_T : S^\alpha \rightarrow M^\beta$ such that

$$e_t \widehat{\theta}_T = \sum_{U \approx T} UC_t^-.$$

By Lemma 2.1, we may assume that U ranges over all tableaux that are row equivalent to T , such that no column of U contains a repeated entry.

Suppose that U, V are α -tableaux. Write $U < V$ if $U \approx V$ and all nodes (a, c) in U with $c > \beta_{a+1}$ contain the symbol a . We say that U is *right-justified*. Now define the following linear combination of α -tableaux (which will belong to M^γ , where γ is the type of V):

$$\widetilde{\theta}_V := \sum_{U < V} UC_t^-.$$

Now recall that T is a semi-standard α -tableau of type β . The analogues of Lemmas 5.2, 5.3, 5.4 and 5.5 hold for $\widehat{\theta}_T$, with the following exception:

$$(16) \quad \widetilde{\theta}_T \psi_a = \widetilde{\theta}_W, \quad \text{if } \hat{T}(a) = a + 1,$$

and not $(h_a - h_{a+1})\widetilde{\theta}_W$. This can be seen by inspecting the original proof. The key point is that there are $\alpha_a - \alpha_{a+1} + 1$ α -tableaux U such that $U \approx T$ and $U\psi_a = W$. However, only one of these U satisfies $U < T$.

Now define the following element of M^β :

$$(17) \quad \mathcal{E}_t := \sum_{\hat{T}(a)=a+1} \left((-1)^{|T|} \prod_{u \in \{R\} \setminus \{T\}} h_u \right) \widetilde{\theta}_T.$$

So T runs over all semi-standard α -tableaux of type β which contain a (unique) symbol $a + 1$ in row a . Lemma 6.5 states that $\widehat{\theta}\psi_{i,r} = 0$, unless $i = a, r = \beta_{a+1} - 1$. In view of the previous paragraph, the analogous result holds for \mathcal{E}_t :

$$\mathcal{E}_t \psi_{i,r} = 0, \quad \text{unless } i = a \text{ and } r = \beta_{a+1} - 1.$$

In view of this, and (4), we see that $e_t\hat{\theta} - h_a\mathcal{E}_t \in S^\beta$, if and only if $(e_t\hat{\theta} - h_a\mathcal{E}_t)\psi_a = 0$. But this equality holds, by Lemma 6.6 and (16). \square

As an example, let $p = 5$ and $\alpha = (4, 3)$ and $\beta = (3^2, 1)$. So $a = 1, b = 3$ and $h_1 = 5, h_2 = 3$. There are two semi-standard α -tableaux of type β :

$$\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} \quad \text{and} \quad \begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 3 \\ \hline 2 & 2 & 2 & \\ \hline \end{array}.$$

corresponding to the semi-standard α -sets $\{1, 2\}$ and $\{1\}$, respectively. Our linear combination $\theta : S_{\mathbb{Z}}^\alpha \rightarrow M_{\mathbb{Z}}^\beta$ is then

$$\theta = \theta \widehat{\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & \\ \hline \end{array}} - 3\theta \widehat{\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 3 \\ \hline 2 & 2 & 2 & \\ \hline \end{array}}.$$

The error term is

$$\mathcal{E}_t = \left(\begin{array}{|c|c|c|} \hline 2 & 1 & 1 & 1 \\ \hline 3 & 2 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 2 & 1 & 1 \\ \hline 2 & 3 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 2 & 1 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} \right) C_t^-.$$

Now consider the following standard α -tableau t and its signed column sum C_t^- :

$$t = \begin{array}{|c|c|c|c|} \hline 1 & 2 & 3 & 4 \\ \hline 5 & 6 & 7 & \\ \hline \end{array}, \quad C_t^- = (1 - (1, 5))(1 - (2, 6))(1 - (3, 7)).$$

We can write $e_t\theta - h_a\mathcal{E}_t$ as a linear combination of α -tabloids:

$$\begin{aligned} e_t\theta - 5\mathcal{E}_t &= \left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 3 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 3 & 2 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 2 & 1 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 2 & 1 & 1 \\ \hline 2 & 3 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 2 & 1 & 1 & 1 \\ \hline 3 & 2 & 2 & \\ \hline \end{array} \right) C_t^- \\ &= \left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 3 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 3 & 1 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 3 & 1 & 1 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 3 & 1 & 1 & 1 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} \right) C_t^- \\ &= \left(\begin{array}{|c|c|c|} \hline 2 & 1 & 1 & 1 \\ \hline 3 & 2 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 2 & 1 & 1 \\ \hline 2 & 3 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 2 & 1 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} \right) C_t^- \\ &= \left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 3 & 1 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} - \begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 3 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} \right) C_t^- \\ &= \left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 3 & 1 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} - \begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 3 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} \right) C_t^- \\ &= \left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 3 & 1 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 2 & 3 \\ \hline 2 & 2 & 1 & \\ \hline \end{array} \right) C_t^- (1 + (2, 3)(6, 7) + (1, 3)(5, 7)). \end{aligned}$$

Now

$$\left(\begin{array}{|c|c|c|} \hline 1 & 1 & 1 & 2 \\ \hline 2 & 2 & 3 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 3 & 1 \\ \hline 2 & 2 & 2 & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 1 & 1 & 2 & 3 \\ \hline 2 & 2 & 1 & \\ \hline \end{array} \right) C_t^- = \left(\begin{array}{|c|c|c|} \hline 3 & 2 & 1 & \\ \hline 4 & 6 & 5 & \\ \hline 7 & & & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 4 & 2 & 1 & \\ \hline 7 & 6 & 5 & \\ \hline 4 & & & \\ \hline \end{array} + \begin{array}{|c|c|c|} \hline 7 & 2 & 1 & \\ \hline 3 & 6 & 5 & \\ \hline 4 & & & \\ \hline \end{array} \right) C_t^- = e_{\begin{array}{|c|c|c|} \hline 3 & 2 & 1 & \\ \hline 4 & 6 & 5 & \\ \hline 7 & & & \\ \hline \end{array}}.$$

Multiplying through by $(1 + (2, 3)(6, 7) + (1, 3)(5, 7))$, we get

$$e_t\theta - 5\mathcal{E}_t = e_{\begin{array}{|c|c|c|} \hline 3 & 2 & 1 & \\ \hline 4 & 6 & 5 & \\ \hline 7 & & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|} \hline 2 & 3 & 1 & \\ \hline 4 & 7 & 5 & \\ \hline 6 & & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 & \\ \hline 4 & 6 & 7 & \\ \hline 5 & & & \\ \hline \end{array}}.$$

We wish to compare this with the expression given by (2). So we use the equalities

$$\begin{aligned} e_{\begin{array}{|c|c|c|} \hline 3 & 2 & 1 & \\ \hline 4 & 6 & 5 & \\ \hline 7 & & & \\ \hline \end{array}} &= e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 & \\ \hline 5 & 6 & 4 & \\ \hline 7 & & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 4 & \\ \hline 5 & 6 & 7 & \\ \hline 3 & & & \\ \hline \end{array}} - e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 & \\ \hline 5 & 6 & 7 & \\ \hline 4 & & & \\ \hline \end{array}} \\ e_{\begin{array}{|c|c|c|} \hline 2 & 3 & 1 & \\ \hline 4 & 7 & 5 & \\ \hline 6 & & & \\ \hline \end{array}} &= e_{\begin{array}{|c|c|c|} \hline 1 & 3 & 2 & \\ \hline 5 & 7 & 4 & \\ \hline 6 & & & \\ \hline \end{array}} + e_{\begin{array}{|c|c|c|} \hline 1 & 3 & 4 & \\ \hline 5 & 7 & 6 & \\ \hline 2 & & & \\ \hline \end{array}} - e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 & \\ \hline 5 & 6 & 7 & \\ \hline 4 & & & \\ \hline \end{array}} \\ e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 & \\ \hline 4 & 6 & 7 & \\ \hline 5 & & & \\ \hline \end{array}} &= -e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 & \\ \hline 5 & 6 & 7 & \\ \hline 4 & & & \\ \hline \end{array}}. \end{aligned}$$

Thus

$$e_t\theta - 5\mathcal{E}_t = e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 5 & 6 & 4 \\ \hline 7 & & \end{array}} + e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 4 \\ \hline 5 & 6 & 7 \\ \hline 3 & & \end{array}} + e_{\begin{array}{|c|c|c|} \hline 1 & 3 & 2 \\ \hline 5 & 7 & 4 \\ \hline 6 & & \end{array}} + e_{\begin{array}{|c|c|c|} \hline 1 & 3 & 4 \\ \hline 5 & 7 & 6 \\ \hline 2 & & \end{array}} - 3e_{\begin{array}{|c|c|c|} \hline 1 & 2 & 3 \\ \hline 5 & 6 & 7 \\ \hline 4 & & \end{array}}.$$

8. ACKNOWLEDGEMENT

Part of this paper was written while Dr. Ellers was visiting the National University of Ireland Maynooth in July/August 2004. The visit was substantially funded by a *New Researcher Award* from the National University of Ireland Maynooth. We gratefully acknowledge this assistance.

REFERENCES

- [1] R. W. Carter, G. Lusztig. On the modular representations of the General Linear and Symmetric groups, *Math. Zeit.* **136** (1974), 193–242.
- [2] R. W. Carter, M. T. J. Payne. On homomorphisms between Weyl modules and Specht modules, *Math. Proc. Camb. Phil. Soc.* **87** (1980), 419–425.
- [3] H. Ellers, J. Murray. Branching rules for Specht modules, *J. Algebra* **307** (2007), 278–286.
- [4] G. D. James. *The Representation theory of the Symmetric Groups* Lecture Notes in Mathematics 682 (Springer-Verlag, 1978).
- [5] A. S. Kleshchev. On Decomposition Numbers and Branching Coefficients for Symmetric and Special Linear Groups, *Proc. London Math. Soc.* **75** (1997), 497–558.
- [6] A. S. Kleshchev. *Linear and Projective Representations of Symmetric Groups* (Cambridge University Press, 2005).
- [7] M. Künzer, G. Nebe. Elementary divisors of Gram matrices of certain Specht modules, *Comm. Alg.* **31** (7) (2003), 3377–3427.
- [8] S. Lyle. Some q -analogs of the Carter-Payne theorem, *J.Reine Angew. Math.* **608** (2007), 93–121.
- [9] G. E. Murphy, A new construction of Young’s seminormal representation of the symmetric groups *J. Algebra* **69** (1981), 287–297.

DEPARTMENT OF MATHEMATICS, ALLEGHENY COLLEGE MEADVILLE, PA 16335, USA
E-mail address: hellers@allegheny.edu

DEPARTMENT OF MATHEMATICS, NATIONAL UNIVERSITY OF IRELAND MAYNOOTH, CO. KILDARE, IRELAND
E-mail address: John.Murray@maths.nuim.ie